



Northeast Fisheries Science Center Reference Document 19-06

# Improving the NEFSC Clam Survey for Atlantic Surfclams and Ocean Quahogs

by Larry Jacobson and Daniel Hennen

May 2019

# Improving the NEFSC Clam Survey for Atlantic Surfclams and Ocean Quahogs

by Larry Jacobson and Daniel Hennen

NOAA Fisheries, Northeast Fisheries Science Center,  
166 Water Street, Woods Hole, MA 02543

**U.S. DEPARTMENT OF COMMERCE**  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Northeast Fisheries Science Center  
Woods Hole, Massachusetts

May 2019

## Northeast Fisheries Science Center Reference Documents

**This series is a secondary scientific series** designed to assure the long-term documentation and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review, and most receive copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

If you do not have Internet access, you may obtain a paper copy of a document by contacting the senior Center author of the desired document. Refer to the title page of the document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate (*i.e.*, non-individualized) authorship, then contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

**Information Quality Act Compliance:** In accordance with section 515 of Public Law 106-554, the Northeast Fisheries Science Center completed both technical and policy reviews for this report. These predissemination reviews are on file at the NEFSC Editorial Office.

This document may be cited as:

Jacobson L, Hennen D . 2019. Improving the NEFSC clam survey for Atlantic surfclams and ocean quahogs. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 19-06; 89 p.  
Available from: <http://www.nefsc.noaa.gov/publications/>

# INTRODUCTION

A working group (WG) consisting of Northeast Fisheries Science Center (NEFSC) and Mid-Atlantic Fishery Management Council (MAFMC) staff, academic partners, and interested persons met 5 times during 2017 to discuss ideas for improving the NEFSC clam survey (Appendix A). The goals were to develop ideas for improving the precision and utility of survey data used in stock assessments and for using survey resources more efficiently. Preliminary ideas were presented to the Mid-Atlantic Fishery Management Council's Scientific and Statistical Committee (SSC) on May 18, 2017, and a final presentation and review of this document by the SSC occurred on March 13, 2018. In addition, there were several meetings within NEFSC between Ecosystems Survey Branch and Population Dynamics Branch staff with leadership participation. This report presents data, analyses, and recommendations for consideration by NEFSC.

The survey targets Atlantic surfclams (*Spisula solidissima*) and ocean quahogs (*Arctica islandica*) in federal waters (US Exclusive Economic Zone or EEZ, 3-200 nautical miles from the coast) along the northeastern coast of the United States between North Carolina and the US-Canada border on Georges Bank (Figure 1). The primary purpose is collection of abundance, biomass, shell length, and meat weight data for both species as well as age data for surfclams which are all used in stock assessments for the EEZ stocks. The hydraulic clam dredge sampling gear is designed to reduce bycatch, and little useful data are collected for other species. Data are routinely collected for additional studies based on sampling requests submitted by NEFSC and external partners prior to each survey. Beginning in 1997, surveys included experimental field work to test sampling gear and estimate capture efficiency and size-selectivity for survey gear. Substantial environmental data (depth and bottom temperatures) have been collected since 1997 but have been little used to date. The fishing industry uses survey results to locate new fishing grounds.

Surfclams are also found in state waters (< 3 nm from shore) and in federal waters of the Gulf of Maine (NEFSC 2017a). However, the NEFSC clam survey does not currently and is not expected to operate in state or Gulf of Maine waters. New Jersey and New York conduct surveys in state waters, but these programs are independent. Federal stock assessments do not include state waters, but information about state resources and fisheries is typically provided in appendices.

Surfclams and ocean quahogs in the US EEZ are managed as single stocks. Their distributions are divided into the northern Georges Bank (GBK) and southern assessment areas for stock assessment purposes (Figure 1). The southern area for both species is divided into 5 regions. The regions (from north to south) are: Southern New England (SNE), Long Island (LI), New Jersey (NJ), Delmarva Peninsula (DMV) and Southern Virginia/North Carolina (SVA). Area and regional boundaries lie along current NEFSC shellfish strata and are based on historical survey, fishery, and stock distribution patterns (Figure 1). This report recommends changes to current survey strata and area/regional boundaries.

Stock size is high (>1 million mt) and exploitation rates are low (<0.01 per year) for the surfclam and ocean quahog stocks as a whole (NEFSC 2016, 2017b). However, stock size is lower and exploitation rates are higher in southern regions, particularly on small productive grounds where the fisheries operate. Fishing grounds tend to be smaller than the 10' square resolution of mandatory logbook reports and are difficult to monitor with the NEFSC clam survey, which is designed to track stock size at larger area or regional scales. Survey and fishery logbook data show that the distributions of surfclams and ocean quahogs and their fisheries shifted northward over the last 30 years. Based on surveys, the proportion of total swept area biomass for surfclams on

GBK changed from about 10% during the 1980s to almost 60% in 2010 (NEFSC 2017b). During the same time period, the proportion of ocean quahog biomass on GBK changed from about 35% during the 1980s to almost 50% (NEFSC 2017c).

Distributions depend on latitude, depth, currents, and temperature for both species. Distributions in southern regions DMV and SVA changed dramatically starting in the 1990s because of relatively warm water conditions and thermal stress (Weinberg 2005). Surfclam habitat begins in the intertidal zone. About 95% of individuals are currently found at depths shallower than 60 m in the north on GBK and shallower than 40 m in the southern DMV region (NEFSC 2017b). Ocean quahogs use deeper habitat with 95% of individuals currently at depths of at least 55 m on GBK and 30-60 m in more southern regions. The distributions of NJ and DMV surfclams shifted into ocean quahog habitat over the last 2 decades making survey and fishery catches with both species more common. The probability of catching both species in the same tow has clearly increased in the NJ region. Habitats for the 2 species are more stable and clearly partitioned by depth in the northern GBK, SNE, and LI regions.

Surfclams may live to at least 30 years compared to ocean quahogs which may live to at least 200 years (Cargnelli et al. 1999a, 1999b). Stock size is more variable for surfclams because they are shorter lived, grow faster, and live in relatively shallow water where environmental variability more strongly effects recruitment, mortality and growth.

The clam survey follows a random stratified design based on NEFSC shellfish strata developed in the late 1970s (Figure 1). The shellfish strata were not designed for clams and are also used for Atlantic sea scallop (*Plactopecten magellanicus*) surveys. Sampling gear and protocols were generally consistent during 1982-2011 with surveys covering both surfclam and ocean quahog habitat every 1-3 years (see Table C6 in NEFSC 2003). In some years, either GBK or the southern assessment area (but not both) were surveyed (NEFSC 2017b, 2017c).

Prior to 2012, a small, lined, hydraulic dredge with a 1.82 m blade was deployed from the NOAA Research Vessel Delaware II. Original protocols specified tows at 1.5 knots for 5 minutes (nominal tow distance 232 m). Beginning in 1997 sensors were used during each dredge tow to measure tow distance directly. The sensor data showed that actual tow distances were much longer than expected (up to 875 m), depth dependent, and varied from survey to survey (Weinberg et al. 2002; NEFSC 2009).

In 2012, the survey was moved to a commercial fishing vessel (F/V *Pursuit*) that carries a more efficient hydraulic dredge with a wider 4 m blade and automatic sorting equipment. Current protocols specify tows at 3 knots for 5 minutes (nominal tow distance 154 m, area swept 617 m<sup>2</sup>). The heavy dredge; faster, free-spooling winches; and improved sensor equipment all reduce uncertainty about tow distance. The commercial dredge and sorting equipment are configured to increase retention of relatively small animals not targeted in the fishery. However, size selectivity is lower for small clams in the current survey than it was historically (NEFSC 2017b, 2017c).

It is important to identify and avoid untowable grounds in the new survey. The commercial dredge used for sampling is highly efficient and robust but difficult to repair at sea. Significant damage from boulders or rough ground would require a trip to port and loss of at least 2-3 days of sampling. A survey might have to be terminated if a dredge were destroyed, and a replacement dredge would be expensive.

Beginning with the introduction of sensor equipment in 1997, surveys included field studies to estimate capture efficiency and size selectivity. Based on these studies, the original survey dredge captured about 23% of large, fully selected surfclams in the path of the dredge while the new commercial dredge captures about 59% (NEFSC 2017b). The original dredge captured

about 19% of large, fully selected ocean quahogs while the new commercial dredge captures about 61% (NEFSC 2017c). These types of experiments are expensive and will be less important in future because capture efficiency and size-selectivity estimates are available for both the original and new surveys. They will be necessary, however, if and when the survey gear changes. The present survey schedule allows for gear and other types of experiments every third year, but there was no need for experimental work during 2017. This report recommends using the time for regular survey sampling.

Decisions about the relative numbers of tows in each stratum during a survey resulted from compromises on optimal allocations for surfclams and optimal allocations for quahogs with ad hoc adjustments to obtain a minimum number of tows in each stratum. Extra tows were added to important regions for special studies and for other purposes. The compromise allocations were far from optimal for either species (see below).

## Key Questions

Discussions centered on the following key and interrelated questions:

1. *Should clam surveys target surfclams and ocean quahogs separately rather than simultaneously?*
2. *Should sampling in poor habitat areas cease, particularly if the 2 species are surveyed separately?*
3. *Should new species-specific stratification schemes be used if the 2 species are surveyed separately?*
4. *Is it feasible to survey the entire stock (GBK plus southern assessment area) for surfclams or ocean quahogs in 1 survey year if the species are separated and sampling area reduced?*
5. *What scheduling options (number, location, and frequency of surveys for both species) should be considered if surveys for the 2 species are separated?*
6. *Can rough ground with risk to equipment damage be avoided?*
7. *Should new strata be constructed from current strata or built from scratch with smaller building blocks?*
8. *How heavily should location and depth information vs. survey catch data be weighted in developing new strata?*
9. *Should new strata schemes with discontinuous strata be considered, or should strata be defined traditionally as single contiguous areas?*
10. *What are the recommended stratification options (method, location, shape, and number of strata) for each species and area?*
11. *Will the recommended changes affect observation and estimation of biological characteristics, such as shell length-weight relationships and growth rates?*
12. *How would potential changes in the clam survey (e.g., lower survey frequency and increased precision) affect management advice and stock assessment modeling?*
13. *How often should future changes in stratification be considered?*
14. *What types of additional research would benefit the clam survey?*

NEFSC staff and the WG addressed each of these questions by using methods described below.

## **MATERIALS AND METHODS**

Two sets of data decisions as well as their analytical approaches are described below. The first set was implemented in the preliminary survey data used by the WG to make recommendations at its final meeting. The remaining decisions were made during or just after the WG's final meeting (see "Data decisions at/after last WG meeting" and "Survey catch density calculations" for details). The latter consist mainly of adjustments to current stratum or regional boundaries and do not materially affect analytical results or recommendations. Where relevant, the data used (preliminary vs. final) are specified below or in the Results section.

### **DMV/SVA**

Clam densities have been very low in the southern and inshore portion of SVA since the 1980s because of warming water, and sampling has been sporadic there (Figure 1). Therefore, strata 1-4, 7, 8, 80, and 81 in SVA were omitted while stratum 5 was combined with DMV to form the new DMV/SVA region (Figures 2-3).

### **Depth Range**

NEFSC clam survey data used in these analyses were collected at depths of 9-80 m, which is the depth range for new recommended survey areas. The innermost stratum boundary in the current survey at 9 m is near the boundary between state and federal waters.

The outermost strata in the current strata set cover 73-110 m, but that entire area has never been fully surveyed. The maximum depth sampled during 1997-2011 was 104 m, the maximum depth sampled by the commercial vessel during 2012-2016 was 75 m, and the maximum depth practical is currently about 80 m. Fishable concentrations may extend out to about 80 m (Dave Wallace, pers. comm.), but quahog catches are low in survey tows at 80 m or deeper. We therefore extended the current strata at 55-73 m out to 80 m and eliminated portions of current strata in deeper water from the survey area (Figures 3-4). For example, the stratum building block identified as "304" is the union of current stratum 3 and the shallow portion of current stratum 4, "708" is used for the union of strata 7 and 8, and so on (Figure 2). Differences in sample density were ignored when deep water strata were combined.

### **Spatial Scale**

It is important to define portions of the survey area to the smallest spatial scale of particular interest (the spatial scale at which it is most important to track abundance) because stratification and other design options are evaluated at this level. The GBK and southern areas were selected because current assessment models allow for separate population dynamics in the 2 areas and because the availability of survey data over time is different for GBK and south assessment areas.

Historically, stock assessments were conducted at smaller regional levels in the south (SNE, LI, NJ, DMV, and SVA, Figure 1), and some analyses at regional or smaller spatial scales are still carried out. Survey designs meant to track regional trends in the south were examined but did not perform well for the southern assessment area as a whole. Stations that could be allocated to precisely track abundance had to be shifted to other strata to adequately track abundance in each individual region, rather than in the southern assessment area overall. In other words, the more important goal of tracking the southern assessment area as a whole conflicted with the goal of tracking individual southern regions. Effects of recommendations on assessment area and regional



survey abundance time series were addressed by the post-stratification and domain analyses described below.

## **Survey Years Used**

The survey data in most analyses were collected during 1997-2016 because sensor data were available to calculate tow distance and because recent data are most relevant to current and future conditions if water temperatures increase as expected. Most of GBK was covered in the new survey during 2013 with the remainder sampled during 2014. Data from both years were combined to form a complete survey for 2013. Similarly, most of SNE was covered during 2013 and 2015 with the rest covered during 2014 and 2016. The SNE data for 2013 and 2014 were combined to form a complete set for 2013. Data for 2015 and 2016 were combined to form a complete set for 2015. Years with complete coverage for GBK were 1986, 1992, 1994, 1997, 1999, 2008, 2011, 2013, and 2016. Years with complete or nearly complete coverage in the southern assessment area were 1982, 1983, 1986, 1989, 1992, 1994, 1997, 1999, 2002, 2005, 2008, 2011, 2012, and 2015.

Water temperatures increased in the southern assessment area and surfclam depth distributions shifted abruptly into deeper water off DMV during the 1990s. Therefore, catch density and station data collected during 1997 and 1999 were omitted in analyses involving only DMV/SVA.

Survey data collected before 1992-1994 were used for “out of sample” (external cross-validation, Picard and Cook 1984) testing as described below. The years 1992-1994 provide a reasonable sample size (2 complete surveys) collected under conditions as similar as possible to 1997-2016, though the lack of sensor data made tow distance impossible to calculate precisely. Years with incomplete sampling for an area were omitted.

## **Building Blocks for New Strata**

Fifteen-minute squares (FMSQ) and current strata (both omitting portions outside 8-80 m and with other modifications described above) were used as “building blocks” in developing new stratification schemes. The decision to use FMSQ was a pragmatic compromise balancing biological and statistical considerations. Larger 20-minute squares were not used because clam densities and ecological characteristics seem to change substantially over distances as large as 20 nm. Ten-minute squares were not used because numbers of tows during 1997-2016 in these smaller areas were usually too small to characterize mean catch with sufficient accuracy. Many FMSQ and some current strata also had low sample sizes, but to a lesser extent. Lack of precision from low sample size was an important consideration in choosing and grouping building blocks to form new strata, in choosing of methods, and in interpreting results.

Regional and area boundaries were modified when using FMSQ because the latter do not fall evenly on current boundary lines. FMSQ entirely within an original region or area were assigned to that original region or area. A square falling on one of the original regional boundaries was assigned based on the amount of area on either side of the original boundary. For example, if a FMSQ lay across the boundary between NJ and LI with 55% of its total area on the NJ side, then the FMSQ would be assigned to the NJ region.



## **Data Decisions at/after Last WG Meeting**

### ***Stratum 73 “Floater”***

The original NEFSC shellfish strata include a small portion of stratum 73 separated from the rest of stratum 73 and surrounded by strata 72 and 74 (Figure 1). For simplicity, and to use contiguous strata building blocks consistently, the floater was assigned to stratum 74 (Figures 3-4).

## **Inconsistent Strata Designations**

In a small number of cases (1.7%), the survey stratum identifier listed on station records did not match the stratum assigned based on the geographic information system (GIS) shapefile for shellfish strata currently used by NEFSC. The discrepancies were caused by small differences in the location of stratum boundaries used historically and in the shapefile to define strata or errors in the recorded stratum. The original stratum designations were replaced by their GIS designation. A small number of tows were omitted entirely because they were outside the survey area based on the GIS designation. This correction had nil impact on results and recommendations but simplified data presentation (Tables 1-5). Adjustments to survey database software should be considered prior to the next clam assessment.

## **GBK/SNE and LI/NJ Regional Boundaries**

The boundary lines for regions and areas lie along stratum boundaries. Based on original NEFSC shellfish strata, the GBK/SNE boundary falls across the southern part of GBK instead of through the Great South Channel as might be expected given geographic and oceanographic patterns (Figure 1.1). Similarly, the current LI/NJ regional boundary is slightly north of Hudson Canyon even though Hudson Canyon is a major feature that strongly effects clam distributions. The original shellfish stratum lines were adjusted so that the GBK/SNE boundary ran through the Great South Channel and the LI/NJ boundary ran through the Hudson Canyon (Figure 3-4). Strata were renumbered where necessary to reflect these changes and to ensure that stratum ID numbers for the GBK and SNE areas were unique. Current stratum 47, in particular, occurs on both sides of the original GBK/SNE boundaries and was renamed such that 471 is in SNE and 472 is on GBK. Any differences in sample density were ignored in redrawing region and area boundary lines.

## **Division of Large Southern Strata by Depth for Ocean Quahogs**

Current strata 5, 9, 13, and 17 cover the 27-46 m depth range in the southern SVA, DMV, and NJ regions (Figure 1). Based on 1997-2017 survey data, surfclams tend to occur across the entire depth range in these strata, but there is a clear tendency for ocean quahogs to occur in the east at depths > 32. We therefore divided strata 5, 9, 13, and 17 at about 32 m in such a way that all or nearly all of the ocean quahog catches occurred east of the dividing lines (Figure 1). Stratum 21 might have been divided as well, but there were small quahog catches near the relatively shallow western border. The decision to divide strata 5, 9, 13, and 17 may increase the precision of quahog surveys because no time will be spent surveying these relatively large areas without any ocean quahog catch and because variance within building blocks variance was reduced.

## *Identifiers for Surfclams and Ocean Quahog Building Blocks*

The building blocks for surfclams and ocean quahogs were different after strata 5, 9, 13, and 17 were divided for ocean quahogs (Figures 3-4). We used the prefix 7 for surfclam and 8 for ocean quahogs to distinguish the 2 sets. For example, current stratum 39 (coded 6039 in the current survey database) becomes building block 739 for surfclams and 839 for ocean quahogs. Current stratum 9 for surfclams becomes 709. The combined 5960 building block in deep water on GBK becomes 75960 for surfclams and 85960 for ocean quahogs. A special convention was used for the quahogs and current strata 5, 9, 13, and 17. For example, the shallow portion of stratum 5 for ocean quahogs becomes building block 89905 while the deep portion becomes building block 805. These conventions had no effect on results or recommendations, but the information is useful in interpreting Tables 1-5 and as general documentation.

## **Nantucket Shoals**

The Nantucket Shoals fishing grounds in federal waters south of Cape Cod, MA, currently supplies a large fraction of the total surfclam catch. The area is not routinely surveyed because of dangerous currents and shallow water. However, it was sampled during 2017 by the Science Center for Marine Fisheries (SCeMFiS) with a small commercial vessel (Powell et al. 2017). The polygon used in designing the 2017 survey may serve as a new stratum if the area is surveyed again (Figure 5).

## *Survey catch density calculations*

Survey data in all analyses were catch numbers rather than catch weight per unit swept area. Catch numbers and weight were highly correlated for surfclams and ocean quahogs, and the same results, decisions, and recommendations would result from using either.

Survey catches were expressed as densities ( $N\ m^{-2}$ ) to account for variation in tow distance, dredge width, and capture efficiency that occurred over time and with replacement of the original survey equipment. Survey catch densities ( $D_t$ ) were computed:

$$D_t = \frac{N_t}{wd_t r}$$

where  $N_t$  is the total catch of clams 40+ mm shell length (SL),  $w$  is the width of the dredge (which changed in 2012),  $d_t$  is the tow distance for each tow based on sensors, and  $r$  is relative capture efficiency (probability of capture for a clam above, below or in front of the path the dredge). The parameters  $r = 0.59/0.23 = 2.57$  for surfclams and  $0.608/0.194 = 3.13$  for ocean quahogs are ratios of capture efficiency estimates and account for fishing power differences in the original (<2012) and current (>2011) surveys. Differences in size selectivity for gear used during the 2 time periods were ignored, but the calculations were restricted to clams 40+ mm shell length (SL).

Survey data for 1997-2016 were pooled in bootstrap and other analyses to increase sample size because spatial patterns were important while interannual variability was not. Interannual differences are clear in the original data, and the variance of the pooled data might greatly exceed the variance among tows in a single survey. Catch data for each year were therefore rescaled to obtain a more realistic level of between tow variance in the pooled data set:

$$d_t = \frac{D_t}{\bar{D}}$$

where  $D_t$  was the catch from tow  $t$  and  $\bar{D}$  was the simple (not stratified) average density in the same spatial domain and year as the tow.

## 1% Rule

The 1% rule was used to reduce the survey area by omitting building blocks in poor habitat areas. A 5% rule was also considered, but it seemed to exclude too much area. The 1% rule guarantees that any building block (FMSQ or current strata) with appreciable densities or abundance are sampled and limits bias in overall swept-area biomass estimates that would be caused by reductions in survey area to less than 1%.

The first step was to calculate the mean density and swept-area abundance (density x area) in each building block for the region or area of interest. The second step was to sort the building blocks by mean density, calculate the cumulative sum of the sorted densities, and then divide by the sum of the densities to compute cumulative proportions. The third step was to repeat these calculations by swept-area abundance. Building blocks with cumulative proportion  $\geq 1\%$  for *either* density or abundance were retained. The swept-area criterion is nearly identical to the density criterion for FMSQ because they are of similar size. The swept-area criterion is more important when using current strata as building blocks because a large, low-density stratum with high abundance could be excluded by using the density criterion alone. The reductions in survey area were smaller when using current strata building blocks.

Preliminary data analyzed before the final WG meeting were scaled before applying the 1% rule. Final analyses used data standardized correctly after applying the 1% rule. These changes affected results, but the overall effect on conclusions and recommendations was nil.

## Optimal Allocation

Random stations were assigned in bootstrap analyses to potential new strata based on a compromise version of Neyman optimal allocation (Neyman 1934). Given a specified total number of random stations (e.g., determined by budget), the Neyman procedure estimates the proportion and number of random stations in each stratum that would minimize the variance of the stratified mean. The calculations are based on stratum area and within-stratum variances (both assumed known precisely). The compromise ensures that at least 2 tows (the minimum number for calculating variance) are assigned to each stratum (stations are removed from strata with the highest allocation so that the total number is unchanged). In practice, optimal allocation calculations are degraded by uncertainty in the stratum variance, particularly if the number of strata is large and number of tows in each stratum is small. Neyman allocation does not include any adjustments for factors that may be important in the survey other than stratum size and variance (e.g., interest in a small area of heavy recruitment).

## Numbers of Stations

Assumptions about the total number of stations in each area were based on survey performance during 2012-2016. During 2012-2016, both GBK and the southern assessment area were covered by the clam survey twice (2 “complete” surveys) with a total of 270 random stations on GBK and 425 random stations in the southern assessment area. Thus  $270/2 = 135$  random

stations per year on GBK and  $425/2 = 212$  random stations per year in the southern assessment area appear feasible. In contrast, the maximum number of random stations per year was 159 for GBK and 186 for the south. Based on these figures, bootstrap and other analyses assumed 150 (about  $(135+159)/2$ ) random stations on GBK and 200 (about  $(212+186)/2$ ) random stations in the southern assessment area. These assumptions had little effect on results and recommendations, but sample size is important in estimating the expected variance in either relative or absolute terms (Appendix B).

The assumed numbers of stations may be optimistic, particularly if funding for clam surveys is reduced. On the other hand, the number of stations could be higher if future surveys target one species at a time and cover a smaller area. Surveys during 2012-2016 included extra days for weather and special field experiments which could also be used for normal survey work but are not included in calculations above. We estimated precision for surveys based on a range of sample sizes to account for potential changes in funding (Appendix B).

## Design Effects

Design effect (DEF) statistics were used to evaluate performance of survey designs. DEF measures the success of a design in reducing the variance of the stratified random mean relative to the variance of a random survey with the same number of tows. The DEF statistic is  $100 \times (1 - \sigma_S^2 / \sigma_R^2)$ , where  $\sigma_S^2$  is the variance of the stratified random mean under the design being tested and  $\sigma_R^2$  is the variance of a random survey with no stratification. DEF can be decomposed into parts by stratification and allocation so that  $DEF = DEF_{Allocation} + DEF_{Stratification}$ . It is important to remember that DEF statistics are estimates that include within-stratum variance estimates that tend to be highly variable. DEF statistics were calculated with the BIOSurvey2 library in R provided by Dr. Stephen Smith (Canada Department of Fisheries and Oceans, emeritus and WG member).

## Historical Comparisons

Effects of separating surveys for the 2 species were evaluated by comparing coefficient of variation (CV) for stratified random means during 1986-2016 (GBK) or 1982-2016 (southern assessment area) using current strata, raw survey data, and BIOSurvey2. The 1% rule was not used, but optimal allocation in these tests used a minimum value of zero stations so that marginal strata could be eliminated entirely. In effect, the test measured effects of both optimal allocation and not sampling marginal habitat. As with other methods used, these calculations understate uncertainty because they assume variance estimates for each stratum in each survey are known.

## Cluster Analysis

Univariate and multivariate cluster analyses were used to group building blocks (FMSQs or current strata) into 2-10 clusters that could potentially be used as new strata. The 2 types of analyses represent different assumptions about accuracy of the mean survey density estimates for building blocks. The univariate method assumes that mean catch estimates for building blocks are accurate while the multivariate analysis accommodates imprecise mean catch estimates by using both location and depth data. Location and depth can be measured accurately although their relationship to spatial patterns in abundance may be inconsistent and likely to change over time.

The univariate procedure tends to form spatially discontinuous clusters (e.g., clusters consisting of building blocks that are scattered across the survey area). In contrast, multivariate procedure tends to group building blocks at adjacent locations and similar depths (e.g., adjacent

FMSQ). The univariate and multivariate analyses are the same when the weights for location and depth in the multivariate case are set to zero.

The univariate procedure was carried out by using the `Ckmeans.1d.dp` library in R with the number of samples in each building block as weights (Wang and Song 2011). It is guaranteed to form clusters that minimize within cluster sums of squares:

$$SSQ = \sum_{c=1}^N \sum_{j=1}^{n_c} n_{c,j} (\hat{d}_{c,j} - \hat{d}_c)^2$$

where  $N$  is the number of clusters specified by the user (an initially arbitrary choice),  $n_c$  is the number of building blocks in cluster  $c$ ,  $n_{c,j}$  is the number of tows for the  $j^{th}$  building block,  $\hat{d}_{c,j}$  the total mean catch, and  $\hat{d}_c = (\sum_j \hat{d}_{scj})/n_c$  is the mean catch in the cluster. This formula is similar to the formula for the variance of a stratified random mean so that minimizing it nearly minimizes the variance of the stratified random mean (when allocation is proportional to area). There is nothing in the formula to reward or penalize proximity in terms of space or depth so that building blocks grouped into the same cluster can be discontinuous and scattered throughout the region.

Multivariate cluster analyses grouped building blocks based on expected catches, position (latitude and longitude), and depth by using the k-means routine in the R stats library (Hartigan and Wong 1979). Multivariate cluster analysis minimized the sum-of-squares ( $SSQ$ ):

$$SSQ = \sum_{c=1}^N \sum_{j=1}^{n_{c,j}} \left[ d'_{s,j}{}^2 + lat'_{s,j}{}^2 + lon'_{s,j}{}^2 + depth'_{s,j}{}^2 \right]$$

where  $lat$  and  $lon$  are positions at the center of a building block and depth is an average calculated over a fine scale grid in each building block. Data on the right hand side of this expression were weighted z-scores. For example,

$$d' = w_d \frac{(d - \hat{d})}{\sigma_d}$$

where,  $w_d$  (default = 1) is a weight that defines the importance of catch density in clustering, and the mean  $\hat{d}$  and standard deviation  $\sigma_d$  for density were computed based on all tows in the analysis. The data were transformed to z-scores because natural differences in scale (e.g., density < 1 and longitude > 100) would overemphasize the variables with larger values. Mean depth, position, and depth are straight-forward for FMSQ but harder to interpret for building blocks based on current strata with complex shapes. The k-means algorithm is a standard approach but not guaranteed to find the best combination of subunits because its starting points and selection process are random. We restarted the algorithm 50 times and kept the best cluster with lowest within-cluster sums of squares.

## Generalized Additive Model (GAM) and Tree models in Multivariate Cluster Analysis

Cluster analysis was also used to group building blocks based on estimated density from Generalized Additive Model GAM and random forest tree models into clusters (Breiman 2001; Wood 2006). Generating population density fields with GAM and random forest tree models was an initial method for increasing the spatial resolution of the data. Artificially increasing the spatial resolution, however, was eventually deemed inappropriate because the resultant population density fields were far more uniform than the actual clam populations. The working group retained the GAM and random forest tree methods for comparison with results using actual survey data, as finding strata that were robust to the patchiness of the population was desirable.

Six GAM models were fit to observed survey catch densities based on location and depth (see below) by using the log link function and assuming the data were from Tweedie distributions to accommodate stations with zero catch. The linear predictors for each model specified in the R programming language were:

<code>~ te(L, k=7) + te(depth, k=7)</code>
<code>~ te(L, k=7) + te(depth, k=7) + as.factor(yr)</code>
<code>~ te(L, k=7) + te(depth, k=7) + te(yr, k=x)</code>
<code>~ te(L, depth, k=14)</code>
<code>~ te(L, depth, k=14) + as.factor(yr)</code>
<code>~ te(L, depth, k=14) + te(yr, k=x)</code>

where  $L$  is location (latitude or longitude), `as.factor(yr)` means different intercept parameters for each year, and  $k$  limits the curvature of univariate [`te(L,k)` `te(depth,k)`] and bivariate [`te(L,depth,k)`] nonlinear terms. Nonlinear terms for year used  $k = \min(7, \text{number of years}-1)$ . Models with year were meant to account for temporal variation in preliminary data induced by applying the 1% rule before rescaling.  $L$  was either latitude or longitude depending on which had the lowest correlation with depth because lack of correlation improves model predictions. The best model was selected based on lowest Akaike information criterion (AIC) and used to make predictions at each point on a fine scale grid (0.01° spacing) within the study area.

Tree models incorporating additional data were used in a similar way. Random forest regression models were fit to observed survey catches based on a depth, latitude, longitude, and environmental and climatological variables (Table 17 in NEFSC 2017c). The fitted tree models were then used to make predictions at each point on the fine scale grid. Missing values in the predictor data were replaced by an imputation routine in the `randomForest` library in R.

Multivariate cluster analyses were used to assign the predicted catch densities from GAM or tree models at each point in the grid into 2-10 clusters, and building block boundaries were ignored (univariate cluster analysis may have performed better). Entire building blocks were assigned to a new stratum based on the most common cluster assigned to the points within it. Eventually, as the number of potential strata increased, one or more strata were lost in the final assignment because the corresponding cluster was not the majority assignment in any of the building blocks. Such cases were abandoned.

## Stratification Options for Bootstrap Tests

A wide range of options with 2-10 new strata based on FMSQ and current strata were compared for both species and regions. Options are identified based on whether current strata or



FMSQ were used, the number of new strata, and the method used to form building blocks. For example, option opt2.3 is based on FMSQ (“opt” means FMSQ building blocks), has 2 new strata, and was developed by using method type 3 (univariate cluster analysis, see table below). Option svd2.3 is analogous but based on current survey strata building blocks (“svd” means current strata as building blocks). The scenarios tested included the current survey design (after applying the 1% rule to current strata) and random assignments of Fifteen Minute Squares (FMSQ) or current strata to new strata. The current stratification scheme with optimal allocation (option opt.quo, 1% rule not applied) was also included.

ID number	Method
1-2	Random
3	Univariate cluster analysis
4	Multivariate cluster analysis
5	GAM model
6	Tree model

## Bootstrapping to Evaluate Stratification Options

Bootstrap techniques were used to evaluate relative performance of stratification options in several types of analyses. Each bootstrap (usually 3000 iterations) involved a simulated survey from an area or region represented by the survey data for 1997-2016 (or earlier years for cross-validation, see below). Tows in future surveys will probably be allocated to strata optimally based on both stratum area and on possibly imprecise variances in abundance estimated from previous survey data. To mimic this pattern in bootstrap analysis, variances were estimated during each bootstrap iteration for each new stratum by using data from a simulated presurvey with the total number of tows allocated to new strata based on relative stratum size (both surveys were simulated in each bootstrap iteration). The total number of tows in presurveys was  $pN$ , where  $N$  is the total number of tows to be carried out in the real survey and  $p$  is a proportion used to reduce the sample size in the presurvey and subsequent accuracy of the variances and allocation ( $p = 1$  in model runs reported here).

It was important in bootstraps to obtain spatially unbiased samples from new strata that were usually composed of portions of 2 or more building blocks originally sampled with different intensities. For example, a new stratum might be composed of 2 building blocks with areas of 10 vs 20 km<sup>2</sup> and with 15 vs. 5 historical tows. Simple random samples from the pooled data give biased bootstrap results because the first building block would be oversampled because of its smaller size and the large number of tows there. To achieve spatially unbiased bootstrap sampling, we first sampled tows from building blocks within a new stratum in proportion to their area based on a multinomial distribution:

$$\vec{n} \sim \text{Multinomial}(N, \vec{p})$$

where  $N$  is the total number of tows in the new stratum after optimal allocation,  $\vec{p}$  is a vector with the proportions of the total area in each old stratum, and  $\vec{n}$  is a random vector holding the number of tows (zero to  $N-1$ ) in each building block ( $\sum \vec{n} = N$ ). The second step was to randomly sample  $n_j$  tows with equal probability from the pooled data for block  $j$ . The stratified random mean, its



CV, design effects (DEF), and other statistics were calculated and stored at the end of each iteration.

Median DEF estimates from each bootstrap iteration were saved and used as the primary performance measure in evaluating stratification options. Distributions of relative errors  $\left(\frac{\hat{x} - x_{true}}{x_{true}}\right)$  where  $x_{true}$  is the “true” value were sometimes used as well. Other statistics including the interquartile range (IQR) for relative errors, median and IQR for  $DEF_{Allocation}$ , and  $DEF_{Stratification}$  and the proportion of DEF statistics  $> 0$  were saved and examined, but DEF was convenient, correlated with the other measures, and seemed to work well. Bias was not considered because all stratification options gave unbiased results if sampling was spatially unbiased. The truth was calculated by calculating the stratified random mean and by using all available data and building blocks as strata (thus, the true value would vary among bootstrap runs with different sets of building blocks).

Bootstrapping was useful, but it is important to point out shortcomings related to sample size and underestimation of variance. In some cases, the number of historical tows in a new stratum or portion of an original stratum was usually low so that the same tows were sampled repeatedly with less variance than would be expected in a real survey. Sample sizes (particularly for FMSQ) were generally low enough that rare large catches were not included. Finally, we used the same data to develop and test stratification options (see exception described below). These shortcomings tend to exaggerate the bootstrap estimates of precision, potential benefits of optimal allocation, and the apparent benefits of increasing numbers of strata. In a real survey, stratum variances are likely to be larger, and the number of optimal strata is likely to be smaller than suggested by the bootstrap results.

## Out of Sample Bootstrap Testing

The performance of a stratification option developed with 1997-2016 data was compared to its performance based on 1992-1994 data. A high correlation between in- and out-of-sample DEF would provide information about how well the option might work in future with new data relative to what might expected based on the original analysis and data. The test was imperfect because stock conditions during 1992-1994 differ from current and future conditions (particularly in the southern assessment area). Nevertheless, the 1992-1994 data were the only available independent data for testing (without reducing the already somewhat sparse 1997-2016 dataset).

## Stability of Clusters

It is possible that cluster results were strongly affected by noise in mean catch-density estimates. A bootstrap-type procedure was used to investigate the stability of clusters and to choose weights for catch density in multivariate analysis.

Stability analysis used station catch densities and building block summary data for each region and species, the information actually used in clustering. These data include the number of tows ( $n_{s,j}$ ), latitude, longitude, and mean depth for each block. The original catch density data (tow-by-tow station records) were pooled into groups defined by their original cluster assignment. For example, if the existing analysis assumed 2 clusters, then all of the catch densities for each station were assigned to one stratum or the other based on the original cluster analysis. In each iteration and for each building block,  $n_{s,j}$  catch densities were sampled with replacements from the pool for the stratum, and the mean of the sample data for each building block was added to a data file with the original latitude, longitude, and depth. The cluster analysis was rerun with these data to assign

each block to a bootstrap cluster, which was recorded. This approach did not fully replicate sampling in actual surveys because catch densities for building blocks within the same original cluster probably have different means and variances. However, there were not enough data to bootstrap catch densities at the individual block level. This procedure was carried out by using the univariate algorithm (effective weight on catch data very large) and the multivariate algorithm with catch weights of 1, 0.5, or 0.25. Weights for location and depth were all set to one.

It is challenging to define and quantify stability in multivariate cluster analysis. For example, imagine FMSQ 1 and 2 were assigned to the cluster 1 with the lowest mean catch density, while FMSQ 3 and 4 were assigned to cluster 2 with the highest mean catch density in the original and first bootstrap analysis. In the second bootstrap analysis, FMSQ 1 and 2 were assigned to the cluster with highest mean catch density, and FMSQ 2 and 3 were assigned to the cluster with the lowest catch density because of a sampling error. The cluster assignments for the 4 FMSQ were 1, 1, 2, 2 in the original and first bootstrap case and 2, 2, 1, 1 in the second bootstrap case. These assignments are different but functionally identical because the first and second FMSQ were always grouped together, as were the third and fourth FMSQ. These issues were surmounted by visual examination of original and bootstrap strata maps (all building blocks assigned to a new stratum plotted in the same color) and by using a root mean square error statistic.

The root mean sum of squares statistic (RMSE) was:

$$RMSE = \sqrt{\frac{\sum_{b=1}^N \sum_{i=1}^{n_R} \sum_{j=1}^{n_R} (O_{i,j} - P_{i,j})^2}{2Nn_R}}$$

where  $N = 3000$  is the number of bootstrap iterations,  $n_R$  is the number of building blocks in the area based on the 1% rule, and  $\mathbf{O}$  and  $\mathbf{P}$  are square ( $n_R \times n_R$ ) symmetrical matrices that record the number of times the  $r^{th}$  and  $c^{th}$  building block occurred in the same cluster based on either the original or  $b^{th}$  bootstrap cluster analysis. Cases where a cluster consisted of a single building block are recorded along the diagonals where  $r = c$ . If the cluster analysis were perfectly stable, then  $\mathbf{O}$  and  $\mathbf{P}$  would always be the same so that  $RMSE = 0$ . The statistic increases as stability is reduced. The statistic is divided by 2 to account approximately for the symmetry of the matrices,  $N$  to account for the number of bootstrap iterations, and  $n_R$  to account for the number of building blocks. Thus, stability is measured as the root mean square number of times that building blocks in the same original cluster were not assigned to the same cluster, per bootstrap iteration and building block. If an error is defined as failure to assign pairs in the original cluster together, then the statistic is the root mean squared error.

The RMSE statistic measures the number of mistakes, not their magnitude or severity. Two stratification schemes in the same region might have the same RMSE statistic even if one scheme involved assignments among adjacent building blocks while the other involved assignments among distant building blocks.

The RMSE statistic is useful for comparing stability of stratification options from cluster analyses with different catch weights within a region, the same species, and the same number of new strata. RMSE results for different numbers of new strata or different areas and species are not comparable.

## Long-term Trends Using Historical Data in New Strata

It was important to evaluate the precision and trends of poststratified means by using new stratification options and historical data. New strata are likely to be fewer, larger, have different boundaries, and cover a smaller total area than the current design. Therefore, current strata that intersect the new survey area will fall entirely or partially within one or more new strata. The conventional formula for stratified means using overlapping sections of the current and new strata is unbiased for poststratification in this case. However, the variance formula for stratified means is biased (variances too small) if sampling rates varied among original (current) strata in the same domain with the new strata (Sarndal et al. 1992). Variances tend to be underestimated because the number of observations in each overlapping section is itself a random variable with variance that is not included in the conventional formula.

We used the `Domain.est` and `post.stratify` functions in `BIOSurvey2` to calculate poststratified means and domain or poststratified variances for surfclams and ocean quahogs in each area and region. Results were compared to stratified, random, mean-catch densities calculated by using the clam survey database used for stock assessment work with current strata. Survey data used in assessments include adjustments not made in the data used elsewhere. Therefore, we used tow data actually employed in the most recent assessments.

## Effects on Stock Assessments and Management Advice

These analyses focus on ocean quahogs because the recommended changes will certainly improve survey data and management advice for surfclam (see Discussion). Data and advice for surfclams would improve because survey frequency would not change, sample density on surfclam habitat would increase, and optimal allocation would be used. The 1% rule will result in negligible negative bias ( $\leq 1\%$ ) in swept-area stock size estimates. Mean survey density will increase but with little or no effect on trends in relative abundance indices unless a substantial portion of the surfclam stock shifts into deep water, (quahog habitat) and is not sampled (see Discussion). Prior distributions for catchability in stock assessment models for surfclams are based on capture efficiency estimates (and availability) and will be unaffected.

The survey frequency for quahogs will change from 1 survey every 3 years to 1 survey every 6 years (see below). Thus, the net effect on survey data and management advice will depend on the tradeoff between gains in precision and losses from reduced survey frequency. This problem was addressed by comparing the relative accuracy of estimated trends in stock size in a simple simulation analysis.

Given the nature of their population dynamics, it is reasonable to approximate ocean quahog stock dynamics with a linear population dynamics model and to approximate the stock assessment model with linear regression. Any changes in the quahog population are likely to be small from year to year. Ocean quahogs grow slowly and experience very low total mortality. Large increases in year class strength may be possible but have probably not occurred during the last century based on preliminary age data (Roger Mann, Virginia Institute of Marine Science, Gloucester Point, Virginia). Any strong year classes that may occur will recruit to the fishable stock slowly over decades so that their effect on stock biomass will be slow and muted. Ocean quahog showed essentially linear abundance trends over the last 40 years, based on a variety of actual stock assessments (Figure 6).

Simulated stock size was  $\mu_{t+1} = \mu_t + \delta\mu_1$ , where the annual increment ( $\delta\mu_1$ ) was constant,  $\mu$  was a parameter and  $t$  was year. Simulated abundance data included lognormal survey errors  $\mu'_t \sim \text{lognormal}(\mu_t, \sigma_s)$ , where  $\mu'_t$  is a stock size estimate and  $\sigma_s$  was the standard deviation of

stratified means from the survey design being considered ( $\sigma_c$  for the current design and  $\sigma_p$  for the proposed design). The standard deviations  $\sigma_s = \sqrt{1 + CV_s^2}$  were based on the CV calculated from errors in stratified mean-catch density during bootstrap analyses using the recommended (new strata and optimal allocation) or current (current strata and mean historical allocation) survey designs with 150 stations on GBK and 200 in the southern assessment areas (Table 6).

The simulated population was “surveyed” by drawing a random  $\mu'_t$  value every 3 years under the previous survey schedule or every 6 years under the recommended survey schedule. After a maximum 50 years of surveys (16 total with 1 every 3 years for surfclams or 10 total with 1 every 6 years for ocean quahogs), a linear regression assessment model was fit to the time series of simulated survey data (without log transforming the data). Estimated trends from the linear models (slope parameters,  $\hat{\delta}$ ) were saved (10,000 iterations) and compared to the true population  $\sigma$  value to compute a relative error  $RE = \frac{(\hat{\delta} - \delta)}{\delta}$  for the estimated rate of population change.

The performance of the survey and assessment model for a scenario was calculated as the standard deviation of the distribution of relative errors in each year. This statistic is a better indicator of survey and assessment model performance than measures of statistical significance because it directly reflects the distance of the estimate from the truth. A lower standard deviation indicates higher precision and accuracy in estimating the true trend in abundance, which is an important objective of stock assessment. Bias was ignored because both survey designs were unbiased, and the mean of the relative errors was always zero.

Sensitivity analyses were used to explore the sensitivity of results to assumptions about the ocean quahog population and survey/assessment assumptions. Sensitivity to the length of the survey time series was tested by varying the length of the simulated survey from 10 to 50 years. Sensitivity to the interval of time between surveys under the proposed design was tested by varying the survey interval from 1 to 12 years, while holding the interval for the current design constant at 3 years. Sensitivity to the variance of estimated abundance in the proposed survey was tested by varying the CV for simulated survey estimates between 0.01 and 1.01 in the proposed survey design, while holding the CV of abundance in the original design constant at the original value. Finally, sensitivity to the direction and magnitude of the true abundance trend was tested by varying  $\delta$  between -0.05 and 0.05.

## Untowable Ground

Untowable ground that might damage the survey dredge was identified with clam survey station records that include location (coordinates at the start of the tow) and information about substrate and gear damage. We focused on Georges Bank where gear damage is most common. Three methods were used.

The first method used substrate data from survey catches during 1980-2011. Data from early surveys were from original paper logs, and data from later surveys were from survey database records. During surveys, data are collected for substrate categories that include “rock” and “boulder.” A score (0, 1 or 2) is assigned to each category based on the watch chief’s estimate of volume. The code 0 meant that no material in the category was observed in the dredge, 1 meant up to 29 bushels, and 2 meant 30 or more bushels. Stations less than one tow distance apart were both assigned the higher score. We designated the location of a tow as untowable ground if the score for “rocks” or “boulders” was a 2.

The second method was a nearest neighbor analysis using data collected on Georges Bank during 1982-2011. The actual process was more complex but, in effect, each tow was categorized

as “towable” or “untowable.” The 4 nearest neighbors in the northeast, southeast, southwest, and northwest quadrants around each “parent” station were identified by using a modified bishop’s move algorithm (Powell et al. 2017). Tows near boundaries were often missing at least 1 of the 4 neighbors. These tows were excluded from further analysis as parents but retained as neighbors. Parent sites with 3 out of 4 neighbors designated “untowable” were deemed to be on untowable bottom.

The last method was based on watch chief comments during 1983-2016 and the SHG (station type, haul type, gear condition) value assigned after each tow. If SHG was greater than 144 (station type random; haul either unrepresentative or aborted because of bad bottom; gear moderately damaged including blade assembly damage, bent bars, knife blade damage and multiple broken nipples) and the damage was determined to be a result of contact with rough bottom by the comments, then the tow location was designated untowable. Tow locations were also designated untowable for those tows in which the watch chief’s comments referenced dredge damage from substrate, large rocks in the tow, or abandonment of the station because of bad ground.

The WG decided to exclude circular areas one nautical mile in diameter centered on the coordinates of each problematic station from the survey area. One nautical mile is (1,852 m), about twice the maximum distance of tows with the old survey dredge (875 m, Weinberg et al. 2002). The area excluded around each station amounts to about 0.785 nm<sup>2</sup> or 2.7 km<sup>2</sup>.

## RESULTS

Surveys areas identified with the 1% rule were smaller than the current survey area by 26%-44% for surfclams and 35%-42% for ocean quahogs (Table 7 and Figures 7-14).

Based on historical analyses, CVs for stratified mean densities in the current survey design were usually similar to what might be expected under random sampling (Figure 15). In contrast, CVs with optimum allocation were always about 50% smaller. DEF calculations for historical surveys (not shown) confirmed these results.  $DEF_{Stratification}$  was usually positive, but  $DEF_{Allocation}$  and DEF were usually small or negative. The benefits of the current survey design were reduced or eliminated by the compromise allocation approach used historically.

Based on bootstrap analyses and DEF with preliminary data, strata options based on FMSQ from univariate cluster analysis performed best for both species in both areas (Figures 16-19). DEF generally increased with the number of strata. The current survey design (with optimal allocation) performed relatively well based on bootstrap results, probably because the numbers of original strata (19 for GBK and 53 in the south) were relatively large. As expected, random assignments of building blocks to new strata performed poorly with median DEF near zero (results not shown).

RMSE scores show that clusters based on FMSQ were relatively stable with catch weights of 0.25 on 0.5 and that univariate clusters, particularly with FMSQ, were unstable (Table 8). The range of RMSE scores for each species and area was wide when using FMSQ as building block indicating that weights had substantial effects on stability. Results for current strata building blocks were mixed with relatively narrow ranges, indicating that weights were less important. Univariate clusters with current strata were relatively stable for GBK surfclams and unstable for GBK ocean quahogs. Strata maps (not shown) confirmed these patterns.

DEF scores from in- and out-of-sample bootstraps with FMSQ were correlated in 2 out of 4 cases (GBK surfclams and GBK ocean quahogs, Figure 20). This correlation implies that for 2 of the 4 area/species combinations, the stratification schemes based on FMSQ worked well for



data not used to generate them. DEF scores with current strata building blocks were correlated in 3 out of 4 cases, suggesting that current strata results may be more stable (GBK surfclams, GBK ocean quahogs, and southern quahogs, Figure 21).

The out-of-sample and stability results with FMSQ and univariate cluster analysis may seem contradictory because performance of FMSQ options was good in some cases (based on in- and out-of-sample bootstrap tests) but always poor in stability tests. The seeming contraction is due to FMSQ with persistent high densities. The univariate procedure tends to consistently (and correctly) cluster the persistent high density FMSQ together in a way that improves precision. Most of the instability, however, was among FMSQ where densities were lower and less certain because of limited data or variability over time. In the lower density cases, random differences in mean density from sampling resulted in substantially different mean catch densities and cluster assignments. Building blocks based on current strata were more stable because they were larger, involved fewer potential combinations, were based on more data, and had relatively certain average catch densities with larger sample sizes in bootstrap analyses.

## Effects on Stock Assessments and Management Advice

Based on simulations, the proposed survey is expected to better detect a trend in ocean quahog abundance than the current survey would under likely future conditions (Figures 22-25). Considering sensitivity results, the standard deviation for relative errors in trend estimates in the proposed survey was less than in the current survey in most cases. The proposed survey was better when the trend in true population was gradual ( $-0.02 < \delta < 0.02$ ). Both surveys performed equally well when the trend in true population was steeper ( $-0.5 < \delta < 0.5$ ). The proposed survey was better over short (10 years) to long (50 years) time periods, when the interval between survey years was less than 9 years, and when the proposed survey was more precise than the current survey.

## Untowable Ground

Method 1 identified 93 problematic locations, method 2 identified 58 locations and method 3 identified 163. There were 314 in problematic locations in of which 208 were unique (Figure 26). The total area excluded from the GBK survey area would be 561 km<sup>2</sup> (163 nm<sup>2</sup>). Most of the excluded areas are on the northern half of GBK where surfclams are common.

## DISCUSSION

Based on the WG decisions during and after its last meeting, multivariate cluster analyses (catch weights of 0.5 and current strata building blocks) and bootstraps were rerun with finalized data. These results together with results based on preliminary data were used to make the final recommendations described below.

## Key Questions and Recommendations

1. *Should clam surveys target surfclams and ocean quahogs separately rather than simultaneously?*
2. *Should sampling of poor habitat areas cease, particularly if the 2 species are surveyed separately?*
3. *Should new species-specific stratification schemes be considered if the 2 species are surveyed separately?*

**Recommendation:** Surfclams and ocean quahogs should be surveyed separately, poor habitat areas should be omitted based on the 1% rule with current strata building blocks, and species-specific stratification schemes should be used to improve precision of data used in stock assessments.

**Rationale:** Historical analyses show that survey precision would improve substantially if stations were allocated optimally for one species at a time with more stations per area of habitat for the target species habitat. Nearly all positive stations for both species were included in new recommended surveys areas that were 26%-44% smaller based on the 1% rule (Table 7 and Figures 7-14). The distribution of surfclams will probably continue moving into deeper water, but there is room for expansion because the recommended survey area for surfclams extends out to 55 m off NJ and DMV/SVA where densities are still low. If surfclams do expand to habitats deeper than 55 m, then they will be caught in surveys targeting ocean quahogs and the survey can be adjusted. Dropping poor habitat areas will generate a negligible negative bias (<1%), but future swept-area biomass estimates will be more accurate because improved precision will offset the bias. Ocean quahogs occur but at very low densities along the 80 m limit of their recommended survey area which is the limit of sampling using the current survey gear. Quahogs recruit and grow slowly and are not likely to establish large dense beds beyond 80 m over the next 2 decades.

4. *Is it feasible to survey the entire stock (GBK plus southern areas) for surfclams or ocean quahogs in one survey year if the species are separated and sampling area reduced?*
5. *What scheduling options (number, location, and frequency of surveys for both species) should be considered if surveys for the 2 species are separated?*

**Recommendation:** According to Survey Branch personnel, it is not feasible to survey the entire stock of either species given available resources and considering survey days lost to weather and equipment problems. Six scheduling options were developed (Table 9). The recommended alternative (Option 5) maintains the current survey frequency for surfclams (2 complete surveys every 6 years) and reduces survey frequency for ocean quahogs from 2 surveys to 1 complete survey every 6 years.

**Rationale:** Both species are at high biomass levels, and overall fishing mortality rates are low because of market limitations. Surfclams older than 30 y and ocean quahogs 200 y are common, so there are no demographic problems likely to result in rapid stock changes. Simulation results suggest that assessment accuracy depends on having precise survey data and reliable information about survey capture efficiency, rather than more frequent but less precise survey data points. It is likely that rapid surfclam declines would be detected at recommended survey frequencies or by using mandatory logbook data which are spatially detailed (reported by trip and 10-minute square) and evaluated annually during routine management. Survey frequency could be adjusted if rapid declines, increased fishing mortality, or any other problems occur.

6. *Can rough ground with risk to equipment damage be avoided?*

**Recommendation:** The rough ground identified in this analysis should be incorporated into station selection algorithms and omitted from the survey area. Strong protocols for dealing with new rough ground during future surveys should be maintained. The database of untowable



grounds should be expanded based on new data from the clam survey, other surveys, and any other data or method available. Untowable ground in the southern region should be identified.

**Rational:** Every effort should be made to avoid grounds where gear damage may occur. The current survey uses a large and expensive commercial dredge mounted on a steep ramp at the end of the vessel, and it is not feasible to make major repairs at sea. Any significant damage to gear at sea will require a return to port, substantial loss of sampling opportunities, and reduced data precision. The database describing untowable grounds should be useful in surveys for species and projects.

7. *Should new strata be constructed from current strata or built from scratch by using smaller building blocks?*
8. *How heavily should location and depth information vs. survey catch data be weighted in developing new strata?*
9. *Should new strata schemes with discontinuous strata be considered or should strata be defined traditionally as single contiguous areas?*
10. *What are the recommended stratification options (method, location, shape, and number of strata) for each species and area?*

**Recommendations:** Based on analyses with final data and all of the WG decisions, the options with 6 strata identified by multivariate cluster analyses using current strata building blocks and catch weights of 0.5 are recommended for both species on GBK and for surfclams in the south (Table 10 and Figures 27-29). The option with 6 strata could also be recommended based on analytical considerations for ocean quahogs in the southern assessment areas (Figure 30). However, one potential stratum for southern ocean quahogs consists of 2 discontinuous groups of building blocks (838 + 83940 in the west and 846 + 84748 in the east). The final recommendation for ocean quahogs in the southern areas separates the discontinuous parts for a total of 7 new strata to reduce the potential impact of operational problems that might reduce the total number of tows and leave the original large stratum partially sampled. These recommendations can be adjusted by Ecosystem Survey Branch to accommodate logistical or other issues not considered here. Modest deviations from these recommendations would not require reanalysis.

## Rationale

Based on median DEF statistics from bootstrap analyses, these recommendations will reduce the variance of stratified random means by 10% for surfclams in the southern area relative to the variance from a simple random design and 25%-35% otherwise (Table 10). In contrast, historical calculations indicate that DEF for the current design is near zero and often negative for both species and areas, although DEF estimates from different types of calculations may be difficult to compare.

The average number of tows allocated to each stratum during bootstrap analysis ranged 2-78 and 6-94 for surfclams in the GBK and in the southern areas (Table 11). The stratum with 2 tows for GBK surfclams could be eliminated entirely if needed. The average number of tows allocated to each stratum ranged 6-56 and 10-55 for ocean quahogs in the GBK and southern areas.

**Rationale:** Judgement is required in making survey design recommendations because uncertainties and limited data make it impossible to identify the best options with certainty. Decisions about contiguity of new strata, building blocks, cluster analysis method, use of survey catch data, and numbers of strata were interrelated, and all were affected by limitations in the

available survey data. Stability analysis with preliminary data showed that multivariate clusters with catch weights of 0.5 and current strata building blocks were relatively stable. Corresponding bootstrap analysis with preliminary data indicated stratification schemes based on these decisions performed reasonably well based on DEF once univariate results were discounted (because lack of stability). Options based on tree and GAM models did not perform particularly well. Out-of-sample bootstrap tests showed that recommended sets worked as expected with “new” data not used to develop the options.

Recommending numbers of strata was a challenging problem. More than 5 strata are seldom required in survey design (Cochran 1977), and the number of strata should be correlated with the amount of information available and inherent variability. The recommended number (6-7 strata) for surfclams and quahogs is larger but more strata may be better, given the constraints on stratum design (contiguity and old stratum building blocks) and given the size and complexity of the clam populations. There was relatively little improvement in DEF or increases in relative errors with more than 6 new strata based on bootstrap analysis with final data (Figures 31-34).

All analytical procedures in this analysis exaggerate the benefits of increasing numbers of strata because the assumed or implicit variance within building blocks and strata were probably too small. Precision (DEF) in survey design depends on trading off benefits from increasing the number of strata (achieving high  $DEF_{Stratification}$ ) and knowing stratum variances well enough to allocate stations to strata accurately (maintaining high  $DEF_{Allocation}$ ). Consider a hypothetically ideal situation where it is easy to develop “good” strata that reflect the spatial distribution of either clam species and reduce within stratum variances. If the variances were known and sample size unlimited, precision could always be improved by increasing the number of strata because the variance in catch density between strata could always be increased. However, stratum variance estimates degrade in real surveys as the number of strata increase because the sample size in each stratum decreases and allocation becomes less accurate. At some point, the gains in  $DEF_{Stratification}$  as strata number increases are offset by loss of  $DEF_{Allocation}$ . The best choice for number of strata is where the gain in  $DEF_{Stratification}$  and loss in  $DEF_{Allocation}$  cancel, and precision is actually lost if more strata are added. Analytical methods used here understate variances and the losses caused by inaccurate allocation so that the point where  $DEF_{Stratification}$  and  $DEF_{Allocation}$  cancel appears shifted in the direction of more strata. The situation is even more complex for clam surveys because the relatively large building blocks based on current strata may make it difficult to match patterns in spatial distributions with stratum boundaries in some cases. Given the misleading nature of such analysis, analysts can choose the number of strata such that the apparent gains in survey performance as additional strata are added are small, as was done here (Figures 31-34).

Small samples tend to more often underestimate than overestimate mean catch within a stratum even though the sampling process is unbiased because the distribution of stratified random means for survey catch are right-skewed (Powell et al. 2017). Variance estimates are more uncertain than mean estimates, and they also tend to be negatively biased with small samples. It may be reasonable to assume that at least 25 stations per stratum are required to estimate variances with adequate precision (Figure 35). If 150 stations can be handled on GBK during one survey, then the number of strata should not exceed  $150/25 = 6$ . Similarly, the number of strata in the southern assessment areas should not exceed  $200/25 = 8$ . If resources for clam surveys were reduced and station numbers fell by 25% then the maximum number of strata would be  $150*0.75/25 = 4.5$  on GBK and  $200*0.75/25 = 6$  in the southern areas. Future reductions in sample size are conceivable if survey funding is limited.

All current NEFSC survey strata are contiguous and based on location and depth without reference to expected catch. This conventional approach is probably necessary in multi-species bottom trawl surveys where spatial patterns in catch vary among species but not in single species NEFSC surveys for invertebrates including clams, sea scallops, and northern shrimp (*Pandalus borealis*). Discontiguous strata are currently used in a relatively small Canadian sea scallop fishery with good results; however, the scallop strata are based largely on habitat information rather than catches (Brown et al. 2012). Discontiguous strata make less sense in very large survey areas such as GBK and the southern area for surfclams and ocean quahogs because distant portions of the same strata are not likely driven by the same environmental patterns. If environmental drivers were better understood, they might be accounted for in stratification procedures. Little is known, however, about the environmental drivers of clam density, and current sampling is too sparse to allow for a thorough exploration of plausible alternatives.

We expected to use FMSQ rather than current strata building blocks to construct new strata that precisely reflected underlying stock distributions, but there were too few survey observations to reliably assign FMSQ to clusters so that cluster results were unstable. The use of current strata building blocks is an inherently conservative approach that has the advantage of helping maintain comparability of survey data collected historically and in the future.

We expected to use univariate cluster analysis based on catch data alone or multivariate cluster analysis with a higher weight on catch, but there was too little catch data (particularly for FMSQ) to weight it heavily without causing instability and discontiguous stratification patterns that were widely scattered and potentially problematic.

The decision to use fewer larger strata enhances potential gains in DEF for allocation because within-stratum variances should be substantially more accurate. It will be important to allocate stations optimally in future surveys, particularly if funding and the total number of tows are reduced. The variability in DEF for recommended options in bootstrap analysis suggests that there may be substantial variability in survey performance and DEF from year to year (Figures 31-34).

Minimizing the variance of the stratified random mean is just one goal in carrying out a survey, and it is possible that future clam surveys will allocate extra tows to important strata (or vice-versa) and make other adjustments where necessary. Similarly, increases to the minimum number of tows per stratum could be considered if searching for new concentrations or monitoring changes in marginal habitat are deemed important.

11. *Will the recommended changes affect observation and estimation of biological characteristics, such as length to weight relationships and growth rates?*

The recommended changes will improve collection of biological data and estimation of biological characteristics because a higher percentage of tows will catch the target species for sampling (more sampling during each survey). Thus, the density of potential samples within the species habitat area will increase. Sampling protocols that call for a fixed number of samples or measurements per stratum will need to be changed, however, because the number of recommended strata is much smaller than the number of current strata.

12. *How would potential changes in the clam survey (e.g., lower survey frequency and increased precision) affect management advice and stock assessment modeling?*

**Recommendation:** The proposed changes in the clam survey are expected to improve management advice and stock assessment modeling for both species in both areas.

**Rationale:** Post-stratification and domain variance statistics indicate that historical trends estimated based on the current and recommended survey designs were similar and that variances under the recommended design were similar or smaller. Post-stratified estimates of mean survey catch density during 1982-2016 were often noticeably higher than estimates from the current survey design because of the 1% rule which excludes poor habitat areas with low catch (Figures 36-43). Swept-area stock-size estimates would be unaffected because the proportional decrease in area was about the same as the increase in mean density.

Simulation analysis shows that more precise survey data will make up for the reduction in survey frequency. Higher survey precision increased a simulated assessment model's ability to measure small changes in stock size for both species and areas. The longer period between surveys for ocean quahog results in larger changes between surveys that are easier to estimate. Rapid declines in ocean quahog stock size are unlikely but might be detected in mandatory logbook data which are spatially detailed (reported by trip and 10-minute square), considered accurate, and evaluated annually. Survey frequency can be adjusted if stock size declines rapidly, fishing mortality increases dramatically, or any other problems occur.

13. *How often should future changes in stratification be considered?*

**Recommendations:** Stations should be allocated to strata based on the best available information, optimal allocation calculations, and other considerations prior to every survey.

Changes might be considered after 10 years (3 surveys for surfclams and 2 for ocean quahogs). However, changes should be avoided unless there is a serious need and substantially more data or new analytical methods for identifying strata. Most of the benefits from recommendations in this report will stem from omitting poor habitat areas from the survey and optimal allocation. Decisions about stratum boundaries were probably less important. Similar to other cases (Smith et al. 1997), a wide range of stratification options appears to work reasonably well for surfclams and ocean quahogs if optimal allocation is employed. The opportunities to reduce survey areas and separate species so that optimal allocation is possible are "low hanging fruit" that will disappear after the recommendations in this report are implemented making substantial future improvements more difficult. A great deal of additional survey data may be required to improve the survey design, but such data will accumulate slowly because clam surveys will be infrequent, particularly for ocean quahogs.

Finally, the expense in making substantial changes to the clam survey should be considered. The recommendations in this report involved a WG consisting of industry representatives, academics from 3 universities, staff from the Greater Atlantic Regional Fisheries Office and Mid-Atlantic Fishery Management Council, several Northeast Fisheries Science Center scientists, and 2 outside experts with statistical and survey expertise funded by the clam industry (Appendix A). Five meetings were required over more than a year of time. External review will require funding as well.

14. *What types of additional research would benefit the clam survey?*

**Recommendations:**

- a. Use data from other surveys bottom type data (e.g., from multibeam sonar) and any other data or methodology available to augment and refine the database used to avoid locations likely to damage sampling gear. Consider using the data to avoid gear damage in other surveys.
- b. Carry out simulation studies to shorten cruise tracks and increase the number of stations that can be occupied in the available time at sea.
- c. Data limitations precluded use of small building blocks (FMSQ) to identify strata. Spatial methods might make better use of available data. Options based on tree and GAM models with location, environmental, and climatological data did not perform well because model estimates of habitat within small areas were patchy and heterogeneous, but the general approaches see promising.

## REFERENCES CITED

- Breiman L. 2001. Random forests. *Machine Learning* 45:5-32.
- Brown CJ, Sameoto JA, Smith SJ. 2012. Multiple methods, maps, and management applications: Purpose made seafloor maps in support of ocean management. *Journal of Sea Research* 72:1-13.
- Cargnelli LM, Griesbach SJ, Packer DB, Weissberger E. 1999a. Essential fish habitat source document: Atlantic surfclam, *Spisula solidissima*, life history and habitat characteristics. NOAA Tech Memo NMFS NE 142; 13 p.
- Cargnelli LM, Griesbach SJ, Packer DB, Weissberger E. 1999b. Essential fish habitat source document: ocean quahog, *Arctica islandica*, life History and habitat characteristics. NOAA Tech Memo NMFS NE 148; 13 p.
- Cochran WG. 1977. Sampling techniques, 3<sup>rd</sup> edition. John Wiley & Sons. NY.
- Hartigan JA, Wong MA. 1979. A K-means clustering algorithm. *Applied Statistics* 28:100-108.
- Neyman J. 1934. On the two different aspects of the representative method: the method of stratified sampling and the method of purposive selection. *Journal of the Royal Statistical Society*, 97(4):558-625.
- Northeast Fisheries Science Center. 2003. Report of the 37th Northeast Regional Stock Assessment Workshop (37th SAW): Stock Assessment Review Committee (SARC) consensus summary of assessments. Northeast Fish. Sci. Cent. Ref. Doc. 03-16; 597 p.
- Northeast Fisheries Science Center. 2009. 48th Northeast Regional Stock Assessment Workshop (48th SAW) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 09-15; 834 p.
- Northeast Fisheries Science Center. 2016. In: 61st Northeast Regional Stock Assessment Workshop (61st SAW) Assessment Summary Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 16-13; 26 p. <http://www.nefsc.noaa.gov/publications/>
- Northeast Fisheries Science Center. 2017a. 63rd Northeast Regional Stock Assessment Workshop (63rd SAW) Assessment Summary Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 17-09; 28 p. <http://www.nefsc.noaa.gov/publications/>
- Northeast Fisheries Science Center. 2017b. 61st Northeast Regional Stock Assessment Workshop (61st SAW) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 17-05; 466 p. <http://www.nefsc.noaa.gov/publications/>
- Northeast Fisheries Science Center. 2017c. 63rd Northeast Regional Stock Assessment Workshop (63rd SAW) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 17-10; 409 p. [www.nefsc.noaa.gov/publications/](http://www.nefsc.noaa.gov/publications/)
- Picard RR, Cook RD. 1984. Cross-validation of regression models. *J Am Stat Assoc* 79(387):575-583.



- Powell EN, Mann R, Ashton-Alcox KA, Kelsey M, Kuykendall KM, Long C. 2017. Can we estimate molluscan abundance and biomass on the continental shelf? *Estuarine and Coastal Science* 198:213-224.
- Powell EN, Kuykendall KM, Moreno P. 2017. The death assemblage as a marker for habitat and an indicator of climate change: Georges Bank, surfclams and ocean quahogs. *Cont. Shelf Res.* 142:14-31.
- Powell EN, Mann R, Kuykendall KM, Long MC, Timbs J. 2018. The “East of Nantucket” survey. Report for the New England Fishery Management Council. Newburyport, MA.
- Sarndal CE, Swenson B, Wretman J. 1992. Model assisted survey sampling. Springer, New York.
- Smith SJ. 1997. Analysis of data from bottom trawl surveys. *Oceanographic Literature Review* 44.7:747-748.
- Wang H, Song M. 2011. Ckmeans.1d.dp: optimal k-means clustering in one dimension by dynamic programming. *The R Journal* 3:29–33.
- Weinberg JR, Rago PJ, Wakefield WW, Keith C. 2002. Estimation of tow distance and spatial heterogeneity using data from inclinometer sensors: an example using a clam survey dredge. *Fisheries Research* 55:49-61.
- Weinberg JR. 2005. Bathymetric shift in the distribution of Atlantic surfclams: response to warmer ocean temperature. *ICES Journal of Marine Science* 62:1444-1453.
- Wood SN. 2006. Generalized additive models. An introduction with R. Chapman and Hall/CRC, New York.



**Table 1. Numbers of positive, zero, and total stations for Atlantic surfclams [*Spisula solidissima*] in the Georges Bank (GBK) area during 1997-2017 that were used in this analysis by building block and original stratum number (final data set reflecting all decisions). Building blocks omitted based on the 1% rule are shaded.<sup>1</sup>**

Area building block, orig. stratum	N positive tows	N zero tows	Total	Area building block, orig. stratum	N positive tows	N zero tows	Total
GBK	330	300	630	75354	16	7	23
763	0	1	1	54	15	7	22
74	0	1	1	55	1	0	1
767	16	13	29	75556	31	5	36
54	1	0	1	55	24	5	29
67	15	13	28	56	6	0	6
768	11	21	32	70	1	0	1
68	11	21	32	75758	33	12	45
769	13	24	37	57	25	12	37
69	12	24	36	58	8	0	8
70	1	0	1	75960	53	43	96
770	27	37	64	59	41	42	83
69	1	0	1	60	12	1	13
70	26	37	63	76162	73	7	80
771	6	16	22	61	57	7	64
71	6	16	22	62	16	0	16
772	12	40	52	76566	8	12	20
72	12	40	52	65	8	12	20
773	7	34	41	747256	9	1	10
69	0	1	1	47	8	1	9
73	7	33	40	55	1	0	1
774	15	27	42				
73	2	3	5				
74	13	24	37				

<sup>1</sup>Note: small numbers of tows from unexpected strata (e.g., one tow from stratum 69 in building block 770) occur because of errors in the original strata designation or because the tow was close to a stratum boundary and small differences between original and current stratum boundaries.

Table 2. Numbers of positive, zero, and total stations for ocean quahogs (*Arctica islandica*) in the Georges Bank (GBK) area during 1997-2017 that were used in this analysis by building block and original stratum number (final data set reflecting all decisions). Building blocks omitted based on the 1% rule are shaded<sup>2</sup>.

Area building block, orig. stratum	N positive tows	N zero tows	Total	Area building block, orig. stratum	N positive tows	N zero tows	Total
GBK	244	386	630	85354	6	17	23
863	1	0	1	54	6	16	22
74	1	0	1	55	0	1	1
867	17	12	29	85556	6	30	36
54	1	0	1	55	5	24	29
67	16	12	28	56	0	6	6
868	24	8	32	70	1	0	1
68	24	8	32	85758	2	43	45
869	32	5	37	57	2	35	37
69	32	4	36	58	0	8	8
70	0	1	1	85960	1	95	96
870	11	53	64	59	1	82	83
69	1	0	1	60	0	13	13
70	10	53	63	86162	1	79	80
871	20	2	22	61	1	63	64
71	20	2	22	62	0	16	16
872	50	2	52	86566	5	15	20
72	50	2	52	65	5	15	20
873	38	3	41	847256	2	8	10
69	1	0	1	47	1	8	9
73	37	3	40	55	1	0	1
874	28	14	42				
73	5	0	5				
74	23	14	37				

<sup>2</sup> Note: small numbers of tows from unexpected strata (e.g., one tow from stratum 69 in building block 770) occur because of errors in the original strata designation or because the tow was close to a stratum boundary and small differences between original and current stratum boundaries.

**Table 3. Numbers of positive, zero, and total stations for Atlantic surfclams [*Spisula solidissima*] in the southern areas during 1997-2017 that were used in this analysis by building block and original stratum number (final data set reflecting all decisions). Building blocks omitted based on the 1% rule are shaded<sup>3</sup>.**

Area building block, orig. stratum	N positive tows	N zero tows	Total	Area building block, orig. stratum	N positive tows	N zero tows	Total	Area building block, orig. stratum	N positive tows	N zero tows	Total
<b>SOUTH</b>	<b>1386</b>	<b>1595</b>	<b>2981</b>	<b>733</b>	<b>51</b>	<b>25</b>	<b>76</b>	<b>793</b>	<b>10</b>	<b>30</b>	<b>40</b>
<b>705</b>	<b>8</b>	<b>41</b>	<b>49</b>	33	51	25	76	33	1	0	1
5	8	40	48	<b>734</b>	<b>87</b>	<b>4</b>	<b>91</b>	93	9	30	39
9	0	1	1	34	87	4	91	<b>794</b>	<b>15</b>	<b>0</b>	<b>15</b>
<b>709</b>	<b>127</b>	<b>237</b>	<b>364</b>	<b>737</b>	<b>18</b>	<b>4</b>	<b>22</b>	94	15	0	15
5	0	1	1	37	18	4	22	<b>795</b>	<b>2</b>	<b>33</b>	<b>35</b>
9	126	234	360	<b>738</b>	<b>44</b>	<b>5</b>	<b>49</b>	95	2	33	35
10	0	1	1	38	43	5	48	<b>796</b>	<b>3</b>	<b>9</b>	<b>12</b>
83	0	1	1	39	1	0	1	96	3	9	12
84	1	0	1	<b>741</b>	<b>45</b>	<b>8</b>	<b>53</b>	<b>71112</b>	<b>16</b>	<b>6</b>	<b>22</b>
<b>710</b>	<b>12</b>	<b>19</b>	<b>31</b>	41	45	8	53	11	16	6	22
6	4	0	4	<b>745</b>	<b>18</b>	<b>18</b>	<b>36</b>	<b>71516</b>	<b>24</b>	<b>7</b>	<b>31</b>
9	0	1	1	45	18	18	36	15	24	7	31
10	8	18	26	<b>746</b>	<b>26</b>	<b>8</b>	<b>34</b>	<b>71920</b>	<b>23</b>	<b>6</b>	<b>29</b>
<b>713</b>	<b>30</b>	<b>116</b>	<b>146</b>	46	26	8	34	19	23	6	29
13	30	116	146	<b>749</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>72324</b>	<b>43</b>	<b>0</b>	<b>43</b>
<b>714</b>	<b>20</b>	<b>29</b>	<b>49</b>	49	1	1	2	23	43	0	43
14	20	29	49	<b>750</b>	<b>3</b>	<b>1</b>	<b>4</b>	<b>72728</b>	<b>26</b>	<b>5</b>	<b>31</b>

<sup>3</sup> Note: small numbers of tows from unexpected strata (e.g., one tow from stratum 69 in building block 770) occur because of errors in the original strata designation or because the tow was close to a stratum boundary and small differences between original and current stratum boundaries.

Table 3, continued. Numbers of positive, zero, and total stations for Atlantic surfclams [*Spisula solidissima*] in the southern areas during 1997-2017 that were used in this analysis by building block and original stratum number (final data set reflecting all decisions). Building blocks omitted based on the 1% rule are shaded

<b>717</b>	<b>27</b>	<b>56</b>	<b>83</b>	49	1	0	1	27	26	5	31
14	0	1	1	50	2	1	3	<b>73132</b>	<b>70</b>	<b>5</b>	<b>75</b>
17	27	54	81	<b>781</b>	<b>6</b>	<b>3</b>	<b>9</b>	26	1	0	1
87	0	1	1	81	6	3	9	27	1	0	1
<b>718</b>	<b>23</b>	<b>6</b>	<b>29</b>	<b>782</b>	<b>6</b>	<b>0</b>	<b>6</b>	30	1	0	1
17	1	0	1	82	6	0	6	31	65	5	70
18	22	6	28	<b>783</b>	<b>7</b>	<b>6</b>	<b>13</b>	32	2	0	2
<b>721</b>	<b>123</b>	<b>226</b>	<b>349</b>	83	7	6	13	<b>73536</b>	<b>53</b>	<b>0</b>	<b>53</b>
21	123	225	348	<b>784</b>	<b>15</b>	<b>13</b>	<b>28</b>	34	3	0	3
25	0	1	1	84	15	13	28	35	48	0	48
<b>722</b>	<b>17</b>	<b>16</b>	<b>33</b>	<b>785</b>	<b>26</b>	<b>21</b>	<b>47</b>	36	1	0	1
22	17	16	33	85	26	21	47	39	1	0	1
<b>725</b>	<b>5</b>	<b>91</b>	<b>96</b>	<b>786</b>	<b>18</b>	<b>7</b>	<b>25</b>	<b>73940</b>	<b>45</b>	<b>1</b>	<b>46</b>
21	0	1	1	86	18	7	25	39	43	1	44
25	4	89	93	<b>787</b>	<b>17</b>	<b>53</b>	<b>70</b>	40	2	0	2
26	1	0	1	21	0	1	1	<b>74344</b>	<b>1</b>	<b>0</b>	<b>1</b>
50	0	1	1	87	17	52	69	39	1	0	1
<b>726</b>	<b>12</b>	<b>7</b>	<b>19</b>	<b>788</b>	<b>18</b>	<b>173</b>	<b>191</b>	<b>75152</b>	<b>3</b>	<b>0</b>	<b>3</b>
26	12	7	19	88	18	172	190	51	3	0	3
<b>729</b>	<b>78</b>	<b>37</b>	<b>115</b>	89	0	1	1	<b>747148</b>	<b>56</b>	<b>3</b>	<b>59</b>
25	10	5	15	<b>789</b>	<b>9</b>	<b>142</b>	<b>151</b>	47	48	3	51
29	67	31	98	89	9	142	151	48	7	0	7
90	1	0	1	<b>790</b>	<b>5</b>	<b>15</b>	<b>20</b>	50	1	0	1
92	0	1	1	90	5	15	20				
<b>730</b>	<b>64</b>	<b>10</b>	<b>74</b>	<b>791</b>	<b>21</b>	<b>36</b>	<b>57</b>				
26	3	3	6	90	5	5	10				
29	0	1	1	91	16	31	47				
30	61	6	67	<b>792</b>	<b>9</b>	<b>56</b>	<b>65</b>				
				29	0	10	10				

**Table 4. Numbers of positive, zero, and total stations for ocean quahogs (*Arctica islandica*) in the southern areas during 1997-2017 that were used in this analysis by building block and original stratum number (final data set reflecting all decisions). Building blocks omitted based on the 1% rule are shaded<sup>4</sup>.**

Area building block, orig. stratum	N positive tows	N zero tows	Total	Area building block, orig. stratum	N positive tows	N zero tows	Total	Area building block, orig. stratum	N positive tows	N zero tows	Total
<b>SOUTH</b>	<b>1457</b>	<b>1524</b>	<b>2981</b>	<b>838</b>	<b>7</b>	<b>42</b>	<b>49</b>	<b>895</b>	<b>12</b>	<b>23</b>	<b>35</b>
<b>805</b>	<b>9</b>	<b>1</b>	<b>10</b>	38	7	41	48	95	12	23	35
5	9	1	10	39	0	1	1	<b>896</b>	<b>12</b>	<b>0</b>	<b>12</b>
<b>809</b>	<b>161</b>	<b>34</b>	<b>195</b>	<b>841</b>	<b>3</b>	<b>50</b>	<b>53</b>	96	12	0	12
5	0	1	1	41	3	50	53	<b>81112</b>	<b>1</b>	<b>21</b>	<b>22</b>
9	160	33	193	<b>845</b>	<b>16</b>	<b>20</b>	<b>36</b>	11	1	21	22
10	1	0	1	45	16	20	36	<b>81516</b>	<b>1</b>	<b>30</b>	<b>31</b>
<b>810</b>	<b>8</b>	<b>23</b>	<b>31</b>	<b>846</b>	<b>8</b>	<b>26</b>	<b>34</b>	15	1	30	31
6	2	2	4	46	8	26	34	<b>81920</b>	<b>0</b>	<b>29</b>	<b>29</b>
9	0	1	1	<b>849</b>	<b>1</b>	<b>1</b>	<b>2</b>	19	0	29	29
10	6	20	26	49	1	1	2	<b>82324</b>	<b>1</b>	<b>42</b>	<b>43</b>
<b>813</b>	<b>38</b>	<b>65</b>	<b>103</b>	<b>850</b>	<b>4</b>	<b>0</b>	<b>4</b>	23	1	42	43
13	38	65	103	49	1	0	1	<b>82728</b>	<b>2</b>	<b>29</b>	<b>31</b>

<sup>4</sup> Note: small numbers of tows from unexpected strata (e.g., one tow from stratum 69 in building block 770) occur because of errors in the original strata designation or because the tow was close to a stratum boundary and small differences between original and current stratum boundaries.

Table 4, continued. Numbers of positive, zero, and total stations for ocean quahogs (*Arctica islandica*) in the southern areas during 1997-2017 that were used in this analysis by building block and original stratum number (final data set reflecting all decisions). Building blocks omitted based on the 1% rule are shaded

<b>814</b>		<b>1</b>	<b>48</b>	<b>49</b>	50	3	0	3	27	2	29	31
	14	1	48	49	<b>881</b>	<b>9</b>	<b>0</b>	<b>9</b>	<b>83132</b>	<b>11</b>	<b>64</b>	<b>75</b>
<b>817</b>		<b>5</b>	<b>54</b>	<b>59</b>	81	9	0	9	26	0	1	1
	14	0	1	1	<b>882</b>	<b>6</b>	<b>0</b>	<b>6</b>	27	1	0	1
	17	5	53	58	82	6	0	6	30	0	1	1
<b>818</b>		<b>1</b>	<b>28</b>	<b>29</b>	<b>883</b>	<b>13</b>	<b>0</b>	<b>13</b>	31	9	61	70
	17	0	1	1	83	13	0	13	32	1	1	2
	18	1	27	28	<b>884</b>	<b>23</b>	<b>5</b>	<b>28</b>	<b>83536</b>	<b>7</b>	<b>46</b>	<b>53</b>
<b>821</b>		<b>173</b>	<b>176</b>	<b>349</b>	84	23	5	28	34	0	3	3
	21	173	175	348	<b>885</b>	<b>45</b>	<b>2</b>	<b>47</b>	35	7	41	48
	25	0	1	1	85	45	2	47	36	0	1	1
<b>822</b>		<b>1</b>	<b>32</b>	<b>33</b>	<b>886</b>	<b>25</b>	<b>0</b>	<b>25</b>	39	0	1	1
	22	1	32	33	86	25	0	25	<b>83940</b>	<b>9</b>	<b>37</b>	<b>46</b>
<b>825</b>		<b>12</b>	<b>84</b>	<b>96</b>	<b>887</b>	<b>68</b>	<b>2</b>	<b>70</b>	39	7	37	44
	21	0	1	1	21	1	0	1	40	2	0	2
	25	11	82	93	87	67	2	69	<b>84344</b>	<b>1</b>	<b>0</b>	<b>1</b>

	26	1	0	1	888	183	8	191	39	1	0	1
	50	0	1	1	88	182	8	190	85152	3	0	3
826		4	15	19	89	1	0	1	51	3	0	3
	26	4	15	19	889	146	5	151	89905	39	0	39
829		6	109	115	89	146	5	151	5	38	0	38
	25	0	15	15	890	14	6	20	9	1	0	1
	29	6	92	98	90	14	6	20	89909	165	4	169
	90	0	1	1	891	26	31	57	9	163	4	167
	92	0	1	1	90	0	10	10	83	1	0	1
830		4	70	74	91	26	21	47	84	1	0	1
	26	1	5	6	892	23	42	65	89913	42	1	43
	29	0	1	1	29	3	7	10	13	42	1	43
	30	3	64	67	92	20	35	55	89917	22	2	24
833		6	70	76	893	1	39	40	17	21	2	23
	33	6	70	76	33	0	1	1	87	1	0	1
834		52	39	91	93	1	38	39	847148	15	44	59
	34	52	39	91	894	7	8	15	47	15	36	51
837		5	17	22	94	7	8	15	48	0	7	7
	37	5	17	22					50	0	1	1



**Table 5. Area and descriptive information for Atlantic surfclam (*Spisula solidissima*) and ocean quahog (*Arctica islandica*) building blocks based on current strata. Areas were taken originally from Northeast Fisheries Science Center (NEFSC) shellfish strata shapefiles and prorated where necessary based on a fine scale grid. GBK = Georges Bank; SNE = Southern New England; LI = Long Island; NJ = New Jersey; DMV/SVA = Delmarva/Southern Virginia.**

Area or region	Building block ID number		Area (km <sup>2</sup> )	Min depth (m)	Max depth (m)	Mean depth (m)	Mean latitude (decimal degrees)	Mean longitude (decimal degrees)
	Surfclam	Quahog						
GBK	763	863	1,052	41.0	80.0	60.7	41.659	-66.900
GBK	767	867	672	18.1	59.9	48.9	41.377	-68.394
GBK	768	868	1,303	11.0	55.0	34.0	41.533	-68.122
GBK	769	869	3,093	21.0	54.0	40.5	41.206	-67.998
GBK	770	870	1,865	37.1	61.2	51.3	40.940	-68.174
GBK	771	871	576	34.0	59.8	49.2	41.995	-67.388
GBK	772	872	1,618	11.1	59.0	37.0	41.808	-67.683
GBK	773	873	1,660	17.0	65.6	40.3	41.525	-67.474
GBK	774	874	1,662	34.5	67.4	52.7	41.663	-67.180
GBK	75354	85354	1,230	42.9	80.0	65.4	41.237	-68.624
GBK	75556	85556	1,476	48.1	80.0	65.7	40.710	-68.741
GBK	75758	85758	781	49.8	80.0	66.6	40.735	-68.224
GBK	75960	85960	2,309	52.8	80.0	68.2	40.908	-67.581
GBK	76162	86162	2,805	41.0	80.0	64.1	41.247	-66.894
GBK	76566	86566	466	37.6	80.0	63.3	41.928	-67.787
GBK	747256	847256	1,052	62.0	80.0	71.6	40.647	-68.981
SNE	737	837	2,263	19.2	55.9	36.6	41.162	-71.242
SNE	738	838	919	43.8	56.5	51.3	40.949	-71.165
SNE	741	841	1,989	22.4	47.8	37.7	41.007	-70.401
SNE	742	842	1,221	44.4	55.6	50.3	40.830	-70.480
SNE	745	845	1,395	20.2	53.7	38.9	40.812	-69.621
SNE	746	846	703	30.3	66.1	51.6	40.705	-69.501
SNE	749	849	761	22.7	52.4	36.9	41.457	-69.571
SNE	750	850	535	37.8	66.6	52.4	41.333	-69.426
SNE	794	894	736	9.0	46.3	24.5	41.275	-71.318
SNE	795	895	953	13.0	31.3	23.3	41.104	-70.263
SNE	796	896	1,680	11.1	44.6	27.3	41.209	-69.615
SNE	73940	83940	3,770	51.0	80.0	64.5	40.678	-71.162
SNE	74344	84344	1,835	53.0	80.0	65.1	40.555	-70.383
SNE	75152	85152	574	49.1	80.0	68.4	41.450	-69.448
SNE	747148	847148	3,178	52.1	80.0	68.2	40.536	-69.499

**Table 5, continued. Area and descriptive information for Atlantic surfclam (*Spisula solidissima*) and ocean quahog (*Arctica islandica*) building blocks based on current strata. Areas were taken originally from Northeast Fisheries Science Center (NEFSC) shellfish strata shapefiles and prorated where necessary based on a fine scale grid. GBK = Georges Bank; SNE = Southern New England; LI = Long Island; NJ = New Jersey; DMV/SVA = Delmarva/Southern Virginia.**

LI	729	829	4,067	25.0	58.4	37.9	40.376	-73.048
LI	730	830	2,398	38.6	78.0	51.4	40.163	-72.797
LI	733	833	1,238	25.0	48.8	38.5	40.802	-72.120
LI	734	834	709	43.8	56.6	50.6	40.666	-72.043
LI	791	891	1,236	9.1	37.5	22.4	40.460	-73.559
LI	792	892	562	9.0	30.9	21.9	40.663	-72.896
LI	793	893	331	9.0	29.1	20.7	40.930	-72.116
LI	73132	83132	3,749	52.0	80.0	65.5	40.048	-72.448
LI	73536	83536	2,775	52.9	80.0	68.1	40.466	-71.815
NJ	717	817	2,411	11.0	52.0	35.6	38.741	-74.353
NJ	718	818	823	42.0	61.0	51.9	38.547	-74.044
NJ	721	821	5,806	18.0	50.4	36.3	39.289	-73.765
NJ	722	822	1,046	42.7	58.8	51.0	39.056	-73.518
NJ	725	825	1,849	25.1	52.5	37.4	39.862	-73.468
NJ	726	826	541	38.0	66.1	50.4	39.846	-73.246
NJ	787	887	1,208	9.0	31.1	16.7	38.985	-74.613
NJ	788	888	1,660	11.6	33.0	21.2	39.383	-74.158
NJ	789	889	1,176	12.1	29.9	22.1	39.884	-73.933
NJ	790	890	337	12.2	37.0	21.5	40.297	-73.891
NJ	71920	81920	1,039	51.3	80.0	64.0	38.460	-73.823
NJ	72324	82324	3,110	47.1	80.0	67.2	38.988	-73.290
NJ	72728	82728	1,728	46.3	80.0	67.6	39.594	-72.920
DMV/SVA	705	---	2,366	15.1	50.0	27.2	36.663	-75.288
DMV/SVA	---	805	507	26.0	50.0	34.9	36.602	-74.964
DMV/SVA	---	89905	1,860	15.1	38.8	25.1	36.680	-75.376
DMV/SVA	709	---	6,496	16.2	58.8	32.6	37.275	-75.126
DMV/SVA	---	809	2,614	26.0	58.8	39.2	37.229	-74.943
DMV/SVA	---	89909	3,882	16.2	38.2	28.1	37.306	-75.249
DMV/SVA	710	810	727	41.5	71.9	53.4	37.146	-74.791
DMV/SVA	713	---	3,940	12.1	51.0	33.6	38.220	-74.639
DMV/SVA	---	813	2,282	23.8	51.0	38.8	38.189	-74.535
DMV/SVA	---	89913	1,659	12.1	39.8	26.5	38.263	-74.782
DMV/SVA	714	814	703	38.2	63.6	51.4	38.078	-74.390
DMV/SVA	781	881	1,241	9.0	25.9	17.3	36.752	-75.770
DMV/SVA	782	882	420	9.0	22.0	15.6	37.056	-75.704
DMV/SVA	783	883	740	9.0	26.1	17.2	37.356	-75.546
DMV/SVA	784	884	1,149	9.0	26.9	16.6	37.771	-75.303

DMVSVA	785	885	1,188	9.0	25.7	16.7	38.244	-74.994
DMVSVA	786	886	675	9.0	34.7	18.3	38.657	-74.942
DMVSVA	71112	81112	1,001	52.0	80.0	64.4	37.319	-74.670
DMVSVA	71516	81516	1,533	45.6	80.0	65.6	37.979	-74.278

**Table 6. Parameters used in simulation analyses to evaluate effects of survey changes on stock assessments and management advice for ocean quahog (*Arctica islandica*) in the Georges Bank (GBK) and southern areas. Coefficient of variation (CV) for survey errors are derived from relative errors = (estimate-truth)/truth in bootstrap analyses assuming either the original survey design and average station allocation or the recommended design with 5 strata and optimal allocation.**

<b>Parameter</b>	<b>GBK</b>		<b>Southern area</b>	
<b>Survey design</b>	<b>Current</b>	<b>Recommended</b>	<b>Current</b>	<b>Recommended</b>
N strata <sup>1</sup>	14	6	40	7
Initial abundance (relative)	0.765	1.25	1.34	1.89
CV for survey errors	0.178	0.103	0.153	0.059
Increment parameter $\delta$	0.01	0.01	0.01	0.01
Survey interval (years)	3	6	3	6

<sup>1</sup>In recommended option (not simulation).

**Table 7. Total area of current survey strata at depths of 9-80 m, area in Georges Bank (GBK) and the southern area retained based on the 1% rule and percent change for Atlantic surfclam [*Spisula solidissima*] and ocean quahog (*Arctica islandica*).**

Area (km <sup>2</sup> )	GBK		South	
	Surfclam	Ocean quahog	Surfclam	Ocean quahog
Total	23,630	23,630	83,290	83,290
Included	17,514	13,652	46,499	54,051
% reduction	26%	42%	44%	35%

Table 8. Root mean squared error (RMSE) statistics measuring stability of potential stratification options with 2-6 new strata based on univariate cluster analysis and multivariate cluster analysis with preliminary data and a range of weights on catch density. Results for Atlantic surfclam [*Spisula solidissima*] and ocean quahog (*Arctica islandica*), areas, numbers of strata, and type of building blocks (current strata vs. fifteen minute squares [FMSQ]) are not comparable. The most stable option in each set (lowest RMSE) is gray, and the least stable (highest RMSE) are bold and italicized.

Surfclams										
Catch Weight	GBK current strata					South current strata				
N new strata	2	3	4	5	6	2	3	4	5	6
0.25	<b>74</b>	<b>66</b>	<b>60</b>	<b>55</b>	39	79	80	81	70	65
0.50	73	65	45	46	40	78	79	80	69	64
1.00	70	62	50	42	<b>42</b>	82	84	82	72	65
Univariate	63	54	51	44	36	<b>93</b>	<b>89</b>	<b>84</b>	<b>78</b>	<b>67</b>
Range	11	12	15	13	6	15	10	4	9	4
Catch Weight	GBK FMSQ					South FMSQ				
0.25	0	8	10	30	7	21	4	19	31	24
0.50	0	20	24	30	12	23	19	27	24	34
1.00	19	18	28	28	23	<b>48</b>	45	42	36	44
Univariate	<b>24</b>	<b>42</b>	<b>31</b>	<b>38</b>	<b>32</b>	30	<b>66</b>	<b>62</b>	<b>60</b>	<b>73</b>
Range	24	35	21	10	25	26	62	43	36	49
Ocean quahogs										
Catch Weight	GBK current strata					South current strata				
N new strata	2	3	4	5	6	2	3	4	5	6
0.25	63	<b>59</b>	37	39	37	116	<b>105</b>	94	<b>87</b>	<b>87</b>
0.50	61	56	46	39	38	116	<b>105</b>	97	<b>87</b>	78
1.00	61	55	<b>52</b>	39	38	116	107	95	84	77
Univariate	<b>64</b>	46	49	<b>42</b>	<b>39</b>	<b>120</b>	101	<b>99</b>	84	76
Range	3	13	16	3	1	4	6	5	3	11
Catch Weight	GBK FMSQ					South FMSQ				
0.25	19	10	1	8	20	18	0	22	23	21
0.50	20	16	1	30	32	17	14	24	17	46
1.00	23	26	9	17	25	35	42	47	68	31
Univariate	<b>36</b>	<b>33</b>	<b>33</b>	<b>32</b>	<b>26</b>	<b>61</b>	<b>63</b>	<b>88</b>	<b>85</b>	<b>77</b>
Range	17	23	32	24	12	44	63	67	68	57

**Table 9. Six options for scheduling Northeast Fisheries Science Center (NEFSC) clam surveys for Atlantic surfclam [*Spisula solidissima*] and ocean quahog (*Arctica islandica*) over an 18 year planning horizon. In the second column, “separate” means that surfclams and ocean quahogs are targeted separately (i.e., during separate years), “test in 3<sup>rd</sup> year” (T) refers to time for gear testing (currently every third year), and “2x surfclams” means that the frequency of surfclam surveys is double the current frequency. “Sn,” “Ss,” “Qs,” and “Qn” refer to surveys for surfclams and ocean quahogs in the northern (Georges Bank) and southern assessment areas. Option 5 (target the 2 species separately, double the survey frequency for surfclams, and halve the frequency for ocean quahogs) is recommended.**

Option	Description	Yr 1	Yr 2	Yr 3	Yr 4	Yr 5	Yr 6	Yr 7	Yr 8	Yr 9	Yr 10	Yr 11	Yr 12	Yr 13	Yr 14	Yr 15	Yr 16	Yr 17	Yr 18	N surveys per year	
																				Surfclam	Quahogs
1	Current	Ss Qs	Sn Qn	T	Ss Qs	Sn Qn	T	Ss Qs	Sn Qn	T	Ss Qs	Sn Qn	T	Ss Qs	Sn Qn	T				0.33	0.33
2	Separate, test in 3rd year	Ss		T	Qs		T	Ss		T	Qs		T	Ss		T				0.20	0.20
			Sn			Qn			Sn			Qn			Sn						
3	Separate, test in 3rd year, 2x surfclams	Ss		T	Ss		T	Qs		T	Ss		T	Ss		T	Qs		T	0.22	0.11
			Sn			Sn			Qn			Sn			Sn			Qn			
4	Separate, no testing	Ss		Qs		Ss		Qs		Ss		Qs		Ss		Qs				0.25	0.25
			Sn		Qn		Sn		Qn		Sn		Qn		Sn		Qn				
5	Separate, no testing, 2x surfclams	Ss		Ss		Qs		Ss		Ss		Qs		Ss		Ss		Qs		0.33	0.17
			Sn		Sn		Qn		Sn		Sn		Qn		Sn		Sn		Qn		
6	Separate, no testing, 3x surfclams	Ss		Ss		Ss		Qs		Ss		Ss		Ss		Qs				0.38	0.13
			Sn		Sn		Sn		Qn		Sn		Sn		Sn		Qn				



**Table 10. Median design efficiency (DEF) statistics for recommended stratification options for the Northeast Fisheries Science Center (NEFSC) clam survey from bootstrap analyses (3000 iterations) for Atlantic surfclam [*Spisula solidissima*] and ocean quahog (*Arctica islandica*). DEF measures the percent reduction in variance of stratified random means relative to the variance from a random design. For example, the recommended option for Georges Bank (GBK) surfclams would be expected to reduce variance by 25% relative to a random design. DEF total = DEF allocation + DEF stratification, where the latter terms are the benefits of optimal allocation and the stratification scheme.**

Species	Area	DEF allocation	DEF stratification	DEF total
Surfclams	GBK	18	7	25
	South	6	4	10
Ocean quahogs	GBK	10	14	25
	South	27	9	35

**Table 11. Bootstrap results for recommended design options in the Northeast Fisheries Science Center (NEFSC) clam survey (stratum level, 3000 iterations) for Atlantic surfclam (*Spisula solidissima*) and ocean quahog (*Arctica islandica*). “Mean variance” is the average within-stratum variances across bootstrap samples, “mean allocation” is the average allocation to each stratum and “perfect allocation” gives the optimal Neyman sample sizes for comparison. There was a total of 150 random stations on GBK and 200 in the south. “Allocation/100 sq km” is the sampling intensity (number of random stations per area). Strata numbers correspond to Figures 27-30.**

Species	Area	N strata	Statistic	New stratum						
				1	2	3	4	5	6	7
Surfclams	GBK	6	Area (sq km)	467	5632	4583	2239	2310	1231	--
			Mean variance	0.29	4.47	23.74	18.55	4.33	5.64	--
			Mean allocation	2	20	78	31	13	6	--
			Perfect allocation	1	34	65	28	14	8	--
			Allocation/100 sq km	0.44	0.35	1.71	1.38	0.54	0.52	--
Surfclams	South	6	Area (sq km)	4733	18076	10104	5305	4221	3300	--
			Mean variance	1.63	5.67	7.17	5.77	4.13	2.88	--
			Mean allocation	9	94	57	27	6	7	--
			Perfect allocation	12	84	53	25	17	11	--
			Allocation/100 sq km	0.19	0.52	0.57	0.50	0.14	0.21	--
Ocean quahogs	GBK	6	Area (sq km)	1663	3760	2806	1866	467	3091	--
			Mean variance	0.89	1.54	1.99	3.69	10.78	7.54	--
			Mean allocation	6	31	27	24	7	56	--
			Perfect allocation	10	29	25	23	10	54	--
			Allocation/100 sq km	0.34	0.81	0.97	1.28	1.54	1.80	--
Ocean quahogs	South	7	Area (sq km)	13037	5828	6837	11364	4689	8414	3881
			Mean variance	1.08	1.86	4.18	11.97	9.26	8.41	87.96
			Mean allocation	17	10	18	55	19	33	48
			Perfect allocation	18	11	19	52	19	33	49
			Allocation/100 sq km	0.13	0.17	0.27	0.48	0.40	0.40	1.24

## APPENDIX A. Working group members, contributors, and attendees.

Individual	Affiliation
Tom Alspach	Sea Watch International
Mike Bergman	NMFS, Northeast Fisheries Science Center
Russ Brown	NMFS, Northeast Fisheries Science Center
Jui-Han Chang	NMFS, Northeast Fisheries Science Center
Nicole Charriere	NMFS, Northeast Fisheries Science Center
Peter Chase	NMFS, Northeast Fisheries Science Center
Toni Chute	NMFS, Northeast Fisheries Science Center
Jessica Coakley	Mid-Atlantic Fishery Management Council
Steve Corriea	Massachusetts Division of Marine Fisheries (retired)
Jonathan Duquette	NMFS, Northeast Fisheries Science Center
Kevin Friedland	NMFS, Northeast Fisheries Science Center
Wendy Gabriel	NMFS, Northeast Fisheries Science Center, Mid-Atlantic Council SSC
Joe Godlewski	NMFS, Northeast Fisheries Science Center
Dvora Hart	NMFS, Northeast Fisheries Science Center
Dan Hennen	NMFS, Northeast Fisheries Science Center
Ed Houde	Center for Environmental Science, University of Maryland
Larry Jacobson	NMFS, Northeast Fisheries Science Center (Chair)
Rob Johnston	NMFS, Northeast Fisheries Science Center
Kelsey Kuykendall	University of Southern Mississippi, SCeMFiS
Paul Kostovick	NMFS, Northeast Fisheries Science Center
Chris Legault	NMFS, Northeast Fisheries Science Center
Chase Long	NMFS, Northeast Fisheries Science Center
Roger Mann	Virginia Institute of Marine Science, SCeMFiS
Nancy McHugh	NMFS, Northeast Fisheries Science Center
Jose Montanez	Mid-Atlantic Fishery Management Council
Daphne Munroe	Haskin Shellfish Research Laboratory, Rutgers University
Vic Nordahl	NMFS, Northeast Fisheries Science Center
Sara Pace	University of Southern Mississippi, SCeMFiS
Doug Potts	NMFS, Greater Atlantic Regional Fisheries Office
Eric Powell	University of Southern Mississippi, SCeMFiS
Paul Rago	NMFS, Northeast Fisheries Science Center (retired), SCeMFiS
Eric Robillard	NMFS, Northeast Fisheries Science Center
Burton Shank	NMFS, Northeast Fisheries Science Center
Michael Simpkins	NMFS, Northeast Fisheries Science Center
Dave Wallace	Wallace and Associates, Ltd.

## APPENDIX B. Effects of sample size (total number of random tows) on survey accuracy for recommended options.

**Recommendation:** If possible, relative Root Mean Square Error (RMSE) statistics should be used to evaluate relationships between accuracy and sample size in Northeast Fisheries Science Center (NEFSC) clam surveys because coefficient of variation (CV) for stratified means may give an overly optimistic impression about survey accuracy (particularly for Georges Bank [GBK] surfclams [*Spisula solidissima*]).

### Methods


Bootstrap analyses using the recommended stratification options (2000 iterations for GBK Atlantic surfclams and 1000 iterations otherwise) were carried out for both species and areas assuming that the total number of random tows was 25, 50, 225, or 250 and with optimal allocation. A relative root mean squared error was used to quantify accuracy:

$$RMSE_N = \frac{\sum_{j=1}^N (x_j - X)^2}{N} / X$$


where  $N$  is the number of random tows,  $X$  is the true survey density, and  $\hat{x}_j$  is the estimated density based on the simulated survey in the  $j^{th}$  bootstrap iteration. We plotted the distribution of CVs [ $SE(x_j)/\hat{x}$ ], the mean CV, and relative RMSE statistics for each case. The distribution central tendencies (median and mean) for CVs are of interest because they are the values encountered in real surveys.

### Results

Results show how survey accuracy improves as sample size increases (Figures Appendix A1-A2). Mean CVs and relative RMSE statistics were always similar, as would be expected in an unbiased survey. Median CVs were noticeably smaller than relative RMSE for surfclams in the GBK and southern areas. These differences reflect the skewed distribution of CV statistics and survey density estimates for surfclams which have more patchy distributions and more variable survey catches. The difference between the median CV and relative RMSE tends to become smaller as sample size increases and the within stratum variance estimates become more accurate.



**Appendix Figure A1.** Bootstrap results (2000 iterations for Georges Bank [GBK] and 1000 iterations for the southern areas) showing relationships between accuracy (average relative root mean squared error, solid line), average survey coefficient of variation (CV) (dash line), and the distribution of survey CVs for Atlantic surfclams (*Spisula solidissima*) in the GBK (top) and southern assessment areas (bottom). The symbol in the middle of the boxplots is at the median, and the block shows the underlying spread. The number of tows assumed in evaluating stratification options (based on historical performance and indicated by arrows) was 150 for GBK and 200 for the southern assessment area.



**Appendix Figure A2. Bootstrap results (2000 iterations for Georges Bank [GBK] and 1000 iterations for the southern areas) showing relationships between accuracy (relative root mean squared error, solid line), average survey coefficient of variation (CV) (dash line), and the distribution of survey CVs for ocean quahogs (*Arctica islandica*) in the GBK (top) and southern assessment areas (bottom). The symbol in the middle of the boxplots is at the mean, and the block shows the underlying spread. The number of tows assumed in evaluating stratification options (based on historical performance and indicated by arrows) was 150 for GBK and 200 for the southern assessment area. Note y-axis scale is different in Figure A2.1.**





Figure 1. Current Northeast Fisheries Science shellfish strata boundaries and area/regions for Atlantic surfclams [*Spisula solidissima*] and ocean quahogs (*Arctica islandica*) established during the late 1970s. The southern area comprises all the regions except Georges Bank.





**Figure 2. Current Northeast Fisheries Science Center shellfish strata boundaries and area/regions for Atlantic surfclams (*Spisula solidissima*) and ocean quahogs (*Arctica islandica*) combined to form building blocks for larger strata.**




Figure 3. Dark lines show new building blocks for surfclams (*Spisula solidissima*). Colors show regional divisions (Georges Bank, Southern New England, Long Island, New Jersey, and Delmarva/Southern Virginia, from north to south). Current Northeast Fisheries Science Center (NEFSC) shellfish strata boundaries are shown as light lines.






Figure 4. Dark lines show new building blocks for quahogs at the southern end of the New Jersey region (blue) and in the Delmarva/Southern Virginia region (green) where Atlantic surfclam (*Spisula solidissima*) and ocean quahog (*Arctica islandica*) building blocks are different. Current Northeast Fisheries Science Center (NEFSC) shellfish strata boundaries are shown as light lines.



**Figure 5. Nantucket Shoals fishing grounds south of Cape Cod, MA.**



**Figure 6. Estimated trends in ocean quahog (*Arctica islandica*) biomass from 7 previous stock assessments and 2 stock assessment models (the older, delay difference KLAMZ model and the currently used statistical catch at age Stock Synthesis model or “SS”).**




**Figure 7. New building blocks (dark lines) and survey area based on the 1% rule for Atlantic surfclams (*Spisula solidissima*) on Georges Bank (GBK). Current survey strata boundaries are shown with the thin line.**




**Figure 8. New building blocks (dark lines) and survey area based on the 1% rule for Atlantic surfclams (*Spisula solidissima*) on Georges Bank (GBK), with survey data from 1997-2016 (rescaled to same annual mean). Hollow symbols show stations with zero catch. Solid symbols show location amount and location of catches.**





**Figure 9. New building blocks (dark lines) and survey area based on the 1% rule for ocean quahogs (*Arctica islandica*) on Georges Bank (GBK). Current survey strata boundaries are shown with the thin line.**



**Figure 10. New building blocks (dark lines) and survey area based on the 1% rule for ocean quahogs (*Arctica islandica*) on Georges Bank (GBK), with survey data from 1997-2016 (rescaled to same annual mean). Hollow symbols show stations with zero catch. Solid symbols show location amount and location of catches.**







Figure 11. New building blocks (dark lines) and survey area based on the 1% rule for surfclams (*Spisula solidissima*) in the southern areas. Current survey strata boundaries are shown with the thin line.




**Figure 12.** New building blocks (lines) and survey area (shaded) based on the 1% rule for surfclams (*Spisula solidissima*) in the southern areas, with survey data from 1997-2016 (rescaled to same annual mean). Hollow symbols show stations with zero catch. Solid symbols show location amount and location of catches.



**Figure 13. New building blocks (dark lines) and survey area based on the 1% rule for ocean quahogs (*Arctica islandica*) in the southern areas. Current survey strata boundaries are shown with the thin line.**



**Figure 14. New building blocks (dark lines) and survey area (shaded area) based on the 1% rule for ocean quahogs (*Arctica islandica*) in the southern areas, with survey data from 1997-2016 (rescaled to same annual mean). Hollow symbols show stations with zero catch. Solid symbols show location amount and location of catches.**



**Figure 15. Historical analysis showing coefficient of variation (CV) for stratified mean catch density in Northeast Fisheries Science Center (NEFSC) clam surveys based on the current design and CVs expected under random sampling and with optimal allocation for Atlantic surfclam (*Spisula solidissima*) and ocean quahog (*Arctica islandica*) in Georges Bank (GBK) and the southern assessment areas. All 3 scenarios use the current stratification scheme.**






Figure 16. Median design efficiency (DEF) results for Atlantic surfclams (*Spisula solidissima*) on Georges Bank (GBK) from bootstrap analysis. Results are shown for 2 types of building blocks (Fifteen-minute squares [FMSQ] vs. Survey Vessel Database System [SVDBS] = current strata), 4 cluster methods, and 2-10 potential new strata (clusters).




Figure 17. Median design efficiency (DEF) results for ocean quahogs (*Arctica islandica*) on Georges Bank (GBK) from bootstrap analysis. Results are shown for 2 types of building blocks (Fifteen-minute squares [FMSQ] vs. Survey Vessel Database System [SVDBS] = current strata), 4 cluster methods, and 2-10 potential new strata (clusters).




Figure 18. Median design efficiency (DEF) results for southern Atlantic surfclams (*Spisula solidissima*) from bootstrap analysis. Results are shown for 2 types of building blocks (Fifteen-minute squares [FMSQ] vs. Survey Vessel Database System [SVDBS] = current strata), 4 cluster methods, and 2-10 potential new strata (clusters).







Figure 19. Median design efficiency (DEF) results for southern ocean quahogs (*Arctica islandica*) from bootstrap analysis. Results are shown for 2 types of building blocks (Fifteen-minute squares [FMSQ] vs. Survey Vessel Database System [SVDBS] = current strata), 4 cluster methods, and 2-10 potential new strata (clusters).




**Figure 20. Median design efficiency (DEF) scores from in-sample (x-axis) and out-of-sample (y-axis) bootstrap analyses for Atlantic surfclam (*Spisula solidissima*) and ocean quahog (*Arctica islandica*) in Georges Bank (GBK) and the southern assessment areas using Fifteen-minute squares (FMSQ) as building blocks and preliminary data.**



**Figure 21. Median design efficiency (DEF) scores from in-sample (x-axis) and out-of-sample (y-axis) bootstrap analyses for Atlantic surfclam (*Spisula solidissima*) and ocean quahog (*Arctica islandica*) in Georges Bank (GBK) and the southern assessment areas using current strata as building blocks and preliminary data.**




**Figure 22. A and B.** Distributions for simulated ocean quahog (*Arctica islandica*) abundance estimates (stratified mean catch density or swept area abundance) in selected years for the original (top) and a preliminary recommended design (bottom). **C.** Annual mean survey abundance, showing the decreasing trend. **D.** Survey samples and 95% confidence bounds taken every 3 years as in the current survey from the distributions depicted in the panel A. **E.** Survey samples taken every 6 years as proposed. Parameters are for the southern area.




**Figure 23. Distributions (n = 10,000) for relative errors in the population trend (slope) parameter  $\frac{(\delta^* - \delta)}{\delta}$  the estimated by weighted linear regression, for the current and preliminarily recommended survey designs.**






**Figure 24. Sensitivity testing for potentially important population parameters for Georges Bank (GBK) ocean quahogs (*Arctica islandica*). The y-axis in each case shows the standard deviation of the distribution of the relative errors shown in Figure 3.16. The dashed line represents the proposed survey, and the solid line represents the current survey.**



**Figure 25. Sensitivity testing for potentially important population parameters for ocean quahogs (*Arctica islandica*) in the southern access areas. The y-axis in each case shows the standard deviation of the distribution of the relative errors shown in Figure 3.16. The dashed line represents the proposed survey, and the solid line represents the current survey.**



**Figure 26. Northeast Fisheries Science Center clam survey stations locations that identify untowable ground. Each location is surrounded by a circle 1 nautical mile in diameter.**




Figure 27. Top: Recommended strata for Georges Bank (GBK) Atlantic surfclams (*Spisula solidissima*) based on cluster and bootstrap analyses with final data, with new stratum ID numbers. Bottom: Recommended strata with survey data used in analyses. Figures are numbered as in Table 11. Each stratum is a different color.




Figure 28. Top: Recommended strata for Georges Bank (GBK) ocean quahogs (*Arctica islandica*) based on cluster and bootstrap analyses with final data, with new stratum ID numbers. Bottom: Recommended strata with survey data used in analyses. Figures are numbered as in Table 11.




Figure 29. Top: Recommended strata for Atlantic surfclams (*Spisula solidissima*) in the southern areas based on cluster and bootstrap analyses with final data, with new stratum ID numbers. Bottom: Recommended strata with survey data used in analyses. Figures are numbered as in Table 11.





Figure 30. Top: Recommended strata for ocean quahogs (*Arctica islandica*) based on cluster and bootstrap analyses with final data, with new stratum ID numbers. Bottom: Recommended strata with survey data used in analyses. Note recommendation to break 1 of the strata identified analytically into 2 new strata based on logistic considerations. Figures are numbered as in Table 11.








Figure 32. Bootstrap results (3000 iterations) for Georges Bank (GBK) ocean quahogs (*Arctica islandica*) and 2-10 new strata (clusters) based on final data. Top left: Relative errors. Top right: design efficiency (*DEF*) (total). Bottom left: *DEF*<sub>Efficiency</sub>. Bottom right: *DEF*<sub>Allocation</sub>. Only positive *DEF* values are shown for clear presentation. Black dots show medians, and notched portion of figures shows approximate 95% confidence intervals for the medians.




Figure 33. Bootstrap results (3000 iterations) for Atlantic surfclams (*Spisula solidissima*) in the southern areas and 2-10 new strata (clusters) based on final data. Top left: Relative errors. Top right: design efficiency ( $DEF$ ) (total). Bottom left:  $DEF_{Efficiency}$ . Bottom right:  $DEF_{Allocation}$ . Only positive  $DEF$  values are shown for clear presentation. Black dots show medians, and notched portion of figures shows approximate 95% confidence intervals for the medians.





Figure 34. Bootstrap results (3000 iterations) for ocean quahogs (*Arctica islandica*) in the southern areas and 2-10 new strata (clusters) based on final data. Top left: Relative errors. Top right: design efficiency ( $DEF$ ) (total). Bottom left:  $DEF_{Efficiency}$ . Bottom right:  $DEF_{Allocation}$ . Only positive  $DEF$  values are shown for clear presentation. Black dots show medians, and notched portion of figures shows approximate 95% confidence intervals for the medians.



**Figure 35. Simulation illustrating relationship between sample size and the precision of estimated means and variances in simple random sampling.** Each panel shows the distribution of 5000 estimated means and variances for samples of the size indicated at the top of the panel (e.g.,  $N = 20$  means 20 tows were sampled 5000 times). The data were for Atlantic surfclams (*Spisula solidissima*) in the southern area during 1997-2016 in building blocks selected based on the 1% rule. The selected data were scaled to a mean on one in each year to avoid exaggerating variance. Vertical lines in each panel show the simulated true population statistic.





Figure 36. Post-stratified means, 95% confidence intervals, and coefficient of variation (CV) based on domain variance calculations and sample size (N tows) for Georges Bank (GBK) Atlantic surfclams (*Spisula solidissima*) calculated using recommended (Domain) and current Stock Assessment Review Committee (SARC) stratification schemes.



**Figure 37. Post-stratified means, 95% confidence intervals and coefficient of variation (CV) based on domain variance calculations and sample size (N tows) for Georges Bank (GBK) ocean quahogs (*Arctica islandica*) calculated by using recommended sampling design (Domain) and current Stock Assessment Review Committee (SARC) stratification schemes.**





Figure 38. Post-stratified means, 95% confidence intervals and coefficient of variation (CV) based on domain variance calculations and sample size (N tows) for Atlantic surfclams (*Spisula solidissima*) in the southern areas calculated by using recommended sampling design (Domain) and current Stock Assessment Review Committee (SARC) stratification schemes.



**Figure 39. Post-stratified means, 95% confidence intervals and coefficient of variation (CV) based on domain variance calculations and sample size (N tows) for ocean quahogs (*Arctica islandica*) in the southern area calculated using recommended sampling design (Domain) and current Stock Assessment Review Committee (SARC) stratification schemes.**







Figure 40. Post-stratified means with 95% confidence intervals, and coefficient of variation (CV) based on domain variance calculations for Atlantic surfclams (*Spisula solidissima*) (top) and ocean quahogs (*Arctica islandica*) (bottom) in the Southern New England (SNE) region calculated by using recommended sampling design (Domain) and current Stock Assessment Review Committee (SARC) stratification schemes.



**Figure 41. Post-stratified means with 95% confidence intervals and coefficient of variation CVs based on domain variance calculations for surfclams (*Spisula solidissima*) (top) and ocean quahogs (*Arctica islandica*) (bottom) in the Long Island (LI) region calculated by using recommended (Domain) and current Stock Assessment Review Committee (SARC) stratification schemes.**





Figure 42. Post-stratified means with 95% confidence intervals and coefficient of variation (CV) based on domain variance calculations for Atlantic surfclams (*Spisula solidissima*) (top) and ocean quahogs (*Arctica islandica*) (bottom) in the New Jersey (NJ) region calculated by using recommended sampling design (Domain) and current Stock Assessment Review Committee (SARC) stratification schemes.



**Figure 43. Post-stratified means with 95% confidence intervals and coefficient of variation (CV) based on domain variance calculations for Atlantic surfclams (*Spisula solidissima*) (top) and ocean quahogs (*Arctica islandica*) (bottom) in the Delmarva/Southern Virginia (DMV/SVA) region calculated by using recommended sampling design (Domain) and current Stock Assessment Review Committee (SARC) stratification schemes.**

# **Procedures for Issuing Manuscripts in the *Northeast Fisheries Science Center Reference Document (CRD) Series***

---

## **Clearance**

All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If any author is not a federal employee, he/she will be required to sign an "NEFSC Release-of-Copyright Form." If your manuscript includes material from another work which has been copyrighted, then you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

For more information, NEFSC authors should see the NEFSC's online publication policy manual, "Manuscript/abstract/webpage preparation, review, and dissemination: NEFSC author's guide to policy, process, and procedure," located in the Publications/Manuscript Review section of the NEFSC intranet page.

## **Organization**

Manuscripts must have an abstract and table of contents, and (if applicable) lists of figures and tables. As much as possible, use traditional scientific manuscript organization for sections: "Introduction," "Study Area" and/or "Experimental Apparatus," "Methods," "Results," "Discussion," "Conclusions," "Acknowledgments," and "Literature/References Cited."

## **Style**

The CRD series is obligated to conform with the style contained in the current edition of the United States Government Printing Office Style Manual. That style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the CSE Style Manual. Manuscripts should be prepared to conform with these style manuals.

The CRD series uses the American Fisheries Society's guides to names of fishes, mollusks, and decapod

crustaceans, the Society for Marine Mammalogy's guide to names of marine mammals, the Biosciences Information Service's guide to serial title abbreviations, and the ISO's (International Standardization Organization) guide to statistical terms.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of cited works. Personal communications must include date, full name, and full mailing address of the contact.

## **Preparation**

Once your document has cleared the review process, the Editorial Office will contact you with publication needs – for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as email attachments or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

## **Production and Distribution**

The Editorial Office will perform a copy-edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once the CRD is ready, the Editorial Office will contact you to review it and submit corrections or changes before the document is posted online.

A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.

---

Research Communications Branch  
Northeast Fisheries Science Center  
National Marine Fisheries Service, NOAA  
166 Water St.  
Woods Hole, MA 02543-1026

**MEDIA  
MAIL**

## **Publications and Reports of the Northeast Fisheries Science Center**

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of living marine resources for the benefit of the nation through their science-based conservation and management and promotion of the health of their environment." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (*e.g.*, anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media. Currently, there are three such media:

*NOAA Technical Memorandum NMFS-NE* -- This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

*Northeast Fisheries Science Center Reference Document* -- This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review and most issues receive copy editing.

*Resource Survey Report* (formerly *Fishermen's Report*) -- This information report is a regularly-issued, quick-turnaround report on the distribution and relative abundance of selected living marine resources as derived from each of the NEFSC's periodic research vessel surveys of the Northeast's continental shelf. This report undergoes internal review, but receives no technical or copy editing.

**TO OBTAIN A COPY** of a *NOAA Technical Memorandum NMFS-NE* or a *Northeast Fisheries Science Center Reference Document*, either contact the NEFSC Editorial Office (166 Water St., Woods Hole, MA 02543-1026; 508-495-2350) or consult the NEFSC webpage on "Reports and Publications" (<http://www.nefsc.noaa.gov/nefsc/publications/>). To access *Resource Survey Report*, consult the Ecosystem Surveys Branch webpage (<http://www.nefsc.noaa.gov/femad/ecosurvey/mainpage/>).

**ANY USE OF TRADE OR BRAND NAMES IN ANY NEFSC PUBLICATION OR REPORT DOES NOT IMPLY ENDORSEMENT.**